

LITTLE THINGS in 3D: robust determination of the circular velocity of dwarf irregular galaxies.

G. Iorio,^{1,2*} F. Fraternali,^{1,3} C. Nipoti,¹ E. Di Teodoro,⁴ J. I. Read⁵ and G. Battaglia^{6,7}

¹*Dipartimento di Fisica e Astronomia, Università di Bologna, Viale Berti Pichat 6/2, I-40127, Bologna, Italy*

²*INAF - Osservatorio Astronomico di Bologna, via Ranzani 1, I-40127, Bologna, Italy*

³*Kapteyn Astronomical Institute, University of Groningen, Landleven 12, 9747 AD Groningen, The Netherlands*

⁴*Research School of Astronomy and Astrophysics - The Australian National University, Canberra, ACT, 2611, Australia*

⁵*Department of Physics, University of Surrey, Guildford, GU2 7XH, Surrey, UK*

⁶*Instituto de Astrofísica de Canarias, calle Via Lactea s/n, E-38205 La Laguna, Tenerife, Spain*

⁷*Universidad de La Laguna, Dpto. Astrofísica, E-38206 La Laguna, Tenerife, Spain*

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

Dwarf Irregular galaxies (dIrrs) are the smallest stellar systems with extended HI discs. The study of the kinematics of such discs is a powerful tool to estimate the total matter distribution at these very small scales. In this work, we study the HI kinematics of 17 galaxies extracted from the ‘Local Irregulars That Trace Luminosity Extremes, The HI Nearby Galaxy Survey’ (LITTLE THINGS). Our approach differs significantly from previous studies in that we directly fit 3D models (two spatial dimensions plus one spectral dimension) using the software ^{3D}BAROLO, fully exploiting the information in the HI datacubes. For each galaxy we derive the geometric parameters of the HI disc (inclination and position angle), the radial distribution of the surface density, the velocity-dispersion (σ_v) profile and the rotation curve. The circular velocity (V_c), which traces directly the galactic potential, is then obtained by correcting the rotation curve for the asymmetric drift. As an initial application, we show that these dIrrs lie on a baryonic Tully-Fisher relation in excellent agreement with that seen on larger scales. The final products of this work are high-quality, ready-to-use kinematic data (V_c and σ_v) that we make publicly available. These can be used to perform dynamical studies and improve our understanding of these low-mass galaxies.

Key words: galaxies: dwarf – galaxies: kinematics – galaxies: ISM – galaxies: structure

1 INTRODUCTION

The HI 21 cm emission line is a powerful tool for studying the dynamics of late-type galaxies since it is typically detected well beyond the optical disc and is not affected by dust extinction. Decades ago, the rise of the radio-interferometers allowed many authors to obtain high-resolution rotation curves for large samples of spiral galaxies (e.g. Bosma 1978a; Bosma et al. 1981; Bosma 1981; van Albada et al. 1985; Begeman 1987). These studies found that the gas rotation remains nearly flat also in the outermost disc, where the visible matter fades and one would expect a nearly Keplerian fall-off of the rotation curve. The flat rotation curves of spirals are one of the most robust indications for the existence of dark matter (DM); gas-rich late-type galaxies are an ideal laboratory for studying the properties of DM halos. In this context, dwarf irregular galaxies (dIrrs) are also studied with great interest. Unlike large spirals, they appear to be dominated by DM down to their very central regions. Thus, the determination of the DM density distribution

is nearly independent of the mass-to-light ratio of the stellar disc (e.g. Casertano & van Gorkom 1991; Côté et al. 2000).

The standard method to extract a galaxy rotation curve is based on the fitting of a tilted-ring model to a 2D velocity field (e.g. Begeman 1987; Schoenmakers et al. 1997). The velocity field is extracted from the HI datacube by deriving the representative velocity of the line profile at each spatial pixel. There are different methods to define a representative velocity, for instance using the intensity-weighted mean (e.g. Rogstad & Shostak 1971), or by fitting functional forms (e.g. single Gaussian, Begeman 1987 or multiple Gaussians, Oh et al. 2015, hereafter O15). The 2D approach has been used in several numerical algorithms such as ROTCUR (Begeman 1987), RESWRI (Schoenmakers et al. 1997), KINEMETRY (Krajnović et al. 2006) and DISKFIT (Spekkens & Sellwood 2007). All of these codes have been very useful to improve our understanding of the kinematics of late-type galaxies. However, working in 2D has a drawback: the results depend on the assumptions made when extracting the velocity field and are also affected by the instrumental resolution (the so-called beam-smearing; Bosma 1978b; Begeman 1987). These problems are especially relevant in

* giuliano.iorio@unibo.it

the study of the kinematics of dIrrs since the HI line profiles can be heavily distorted by both non-circular motions and noise. As a consequence, the extraction of the velocity field can be challenging (see e.g. [Oh et al. 2008](#)). Furthermore, dIrrs are often observed with a limited number of resolution elements because of their limited extension on the sky. The effect of beam smearing can also make the estimate of both the rotation curve and the inclination of the disc uncertain ([Begeman 1987](#)).

The pioneering work of [Swaters \(1999\)](#) showed that these problems can be solved with an alternative approach: the properties of the HI disc can be retrieved with a direct modelling of the 3D datacube (two spatial axes plus one spectral axis) without explicitly extracting velocity fields. In practice, a 3D method consists of a data-model comparison of n_{ch} maps (where n_{ch} is the number of spectral channels) instead of the single map represented by the velocity field. The best advantage of this approach is that the datacube models are convolved with the instrumental response, so the final results are not affected by the beam smearing. [Swaters \(1999\)](#), [Gentile et al. \(2004\)](#), [Lelli et al. \(2010\)](#) and [Lelli et al. \(2014b\)](#) used a 3D visual comparison between datacubes and model cubes to correct and improve the results obtained with the classical 2D methods. However, a by-eye inspection of the data is time intensive and subjective. Modern software like TiRiFiC ([Józsa et al. 2007](#)) and ^{3D}BAROLO (hereafter 3DB; [Di Teodoro & Fraternali 2015](#)) can perform a full 3D numerical minimisation on the whole datacube.

In this paper, we exploit the power of the 3D approach by applying 3DB to a sample of 17 dIrrs taken from the LITTLE THINGS (hereafter LT) sample ([Hunter et al. 2012](#)). 3DB has been extensively tested on mock HI datacubes corresponding to low mass isolated dIrrs. [Read et al. \(2016c\)](#) show that the rotation curve is well-recovered even for a star-bursting dwarf, provided that the inclination of the HI disc is higher than 40° .

The final products of our analysis are ready-to-use rotation curves¹ corrected for the instrumental effects (see Sec. 3.2) and for the asymmetric drift (see Sec. 4.3.1). The rotation curves can be used to study the dynamics of dIrrs (e.g. [van Eymeren et al. 2009b](#); [Read et al. 2016a,c](#)) or to extend the study of the scaling relations on small scales (e.g. [Brook et al. 2016a](#); [Lelli et al. 2016a](#)). An unbiased estimate of the rotation curves of dIrrs is also essential to shed light on the well-known cosmological tensions on small scale: the long debated ‘cusp-core problem’ ([de Blok & Bosma 2002a](#); [Oh et al. 2011](#); [Read et al. 2016b](#)), the ‘missing-satellites problem’ ([Klypin et al. 1999](#); [Read et al. 2016a](#)) and the ‘too-big-to-fail problem’ ([Read et al. 2006](#); [Boylan-Kolchin et al. 2011](#); [Read et al. 2016a](#)). These “problems” are discrepancies between the properties of DM halos predicted by cosmological simulations and those inferred from observations. In this context it is therefore crucial to obtain reliable estimates of the DM distribution in real galaxies.

In this work, we focus in particular on the asymmetric-drift correction (see Sec. 4.3.1). This correction term is usually applied without considering the errors made in its calculation, but these can be very large and should be taken into account. The propagation of the asymmetric-drift errors has a great impact on the determination of the circular velocity of galaxies in which the contribution of the gas velocity dispersion is important (see for example DDO 210 in Sec. 5). We developed a method to calculate and propagate the asymmetric-drift uncertainties: as a result, the quoted errors on the

final corrected rotation curves are a robust description of the real uncertainties.

The paper is organised as follows. In Section 2, we illustrate the sample and the data used in this work. In Section 3, we briefly introduce the tilted-ring model and 3DB. In Section 4, we describe in detail the analysis applied to our data. Section 5 shows the results of our analysis for each galaxy in our sample. In Section 6, we look at whether the low mass dIrrs that we study here lie on the baryonic Tully-Fisher relation. In Section 7, we discuss the assumptions made to derive the rotation curves and we compare our results with previous work. A summary is given in Section 8.

2 SAMPLE AND DATA

2.1 Sample selection

The galaxies used in this work represent a sub-sample of the galaxies of the LT survey ([Hunter et al. 2012](#)) which is a sample of 37 dIrrs and 4 Blue Compact Dwarfs (BCDs) located in the Local Volume within 11 Mpc from the Milky Way. LT combines archival and new HI observations taken with the Very Large Array (VLA) to obtain a very high spatial and spectral resolution data set (see Table 1). The objects in the LT sample have been chosen to be both isolated and representative of the full range of dIrr properties ([Hunter et al. 2012](#)).

We built our sample selecting dIrrs that are well-suited to study the circular motion of the HI discs. To this end, we excluded from our selection all the objects seen at low inclination angles and the 4 BCDs. The inclination angle i is defined as the angle between the plane of the disc and the line of sight such that $i = 90^\circ$ for an edge-on galaxy and $i = 0^\circ$ for a face-on galaxy. The estimate of i in nearly face-on galaxies ($i < 40^\circ$) is extremely difficult both from the kinematic fit ([Begeman 1987](#)) and from the analysis of the HI map ([Read et al. 2016c](#)). Moreover, for $i < 40^\circ$, relatively small errors on i have a great impact on the deprojection of the observed rotational velocity (see Eq. 1). As a consequence, the final rotation curves of these nearly face-on galaxies could be biased and unreliable with both 2D ([Oman et al. 2016](#)) and 3D methods ([Read et al. 2016c](#)). Among the objects with $i > 40^\circ$, we selected 17 dIrrs, about half of the original LT sample.

We checked, using a Kolmogorov-Smirnov (KS) test, that our sub-sample preserves the statistical distribution of the galactic properties such as distance (KS p-value 0.96), absolute magnitude (KS p-value 0.99), star formation rate density (SFRD, KS p-value 0.92) and baryonic mass (KS p-value 0.99). Because of our rejection criterion, the average i in our sub-sample (58°) is higher than the one measured in the LT sample (50°). Figure 1 shows the distributions of the distances and of the absolute magnitudes of our sample and the original LT sample.

It is worth pointing out that our rejection criterion is based on i estimated for the stellar disc from the V-band photometry ([Hunter & Elmegreen 2006](#)). When we re-estimated i from the HI datacubes (see Sec. 4.2), we found that four galaxies (DDO 47, DDO 50, DDO 53, and DDO 133) have an average i lower than 40° (see Table 2). We did not exclude these galaxies from our sample, but our results for these objects should be treated with caution. In particular the formal errors on the velocities and on i could underestimate the real uncertainties (see notes on the individual galaxies in Sec. 5).

In conclusion, our sample comprises 17 objects covering the stellar mass range $5 \times 10^5 \lesssim M_*/M_\odot \lesssim 10^8$, with a mean SFRD

¹ The final rotation curves can be downloaded from <http://www.filippofraternali.com/styled-9/index.html>




Figure 1. Distribution of the Galactocentric distances (left-hand panel) and of the absolute magnitudes (right-hand panel) for the sample of galaxies studied in this work (red histograms) and for the whole LT sample (blue dashed histograms). Data from [Hunter et al. \(2012\)](#) and references therein.

of about $0.006 \text{ M}_{\odot} \text{ yr}^{-1} \text{ kpc}^{-2}$ and a mean specific SFRD (SSFRD) of about $2.26 \times 10^{-10} \text{ yr}^{-1} \text{ kpc}^{-2}$ (data from [Hunter et al. 2012](#) and references therein). For comparison the SFRD and the SSFRD of the complete LT sample are $0.007 \text{ M}_{\odot} \text{ yr}^{-1} \text{ kpc}^{-2}$ and $2.95 \times 10^{-10} \text{ yr}^{-1} \text{ kpc}^{-2}$ respectively ([Hunter et al. 2012](#) and references therein).

2.2 The Data

The HI datacubes were taken from the the publicly available archive of the LT survey². For each galaxy, LT provides two datacubes that differ in the weight scheme used to reduce the raw data: the natural datacube offers the best signal-to-noise ratio, the robust datacube gives the best angular resolution. We preferred the natural datacubes for two main reasons: (i) an high signal-to-noise ratio allows us to have enough sensitivity to extend the study of the kinematics at large radii; (ii) 3D methods are not dramatically influenced by the resolution of the data (see Sec. 3). In some cases (DDO 50, DDO 154, NGC 1569, NGC 2366, UGC 8508 and WLM) the galaxies are so extended in the sky that we further smoothed the datacube maintaining a good spatial resolution. Instead, we have chosen the robust datacube when the natural datacube gives too few resolution elements to sample the HI disc (5 cases, see Tab. 1). The mean spatial resolution is about 240 pc ranging from about 60 pc (DDO 210) to about 460 pc (DDO 87). These data are summarised in Cols. 4-6 of Table 1. The rotation curves of all the galaxies in our sample have been already derived with the classical 2D approach in O15. The masses and the surface-density profiles of the stellar discs for all the galaxies can be found in [Zhang et al. \(2012\)](#), except for DDO 47 for which they can be found in [Makarova \(1999\)](#) and in [Walter & Brinks \(2001\)](#).

² <https://science.nrao.edu/science/surveys/littlethings>

3 THE METHOD

3.1 Tilted-ring model

Since [Rogstad et al. \(1974\)](#), the most common approach to study the kinematics of the HI discs is the so-called ‘tilted-ring model’. In practice the rotating disc is broken into a series of independent circular rings with radius R , each with its kinematic and geometric properties. The projection of each ring on the sky is an elliptical ring with semi-major axis R and semi-minor axis $R \cos i$. Throughout the paper R indicates both the radius of the circular ring and the semi-major axis of its projection. For a given projected ring, the projected velocity along the line of sight (V_{los}) is

$$V_{\text{los}} = V_{\text{sys}} + V_{\text{rot}}(R) \cos \theta \sin i, \quad (1)$$

where V_{sys} is the systemic velocity of the galaxy, V_{rot} is the rotation velocity of the gas at radius R , θ is the azimuthal angle of the rings in the plane of the galaxy (related to i , to the galaxy centre and to the position angle; Eq. 2a and 2b in [Begeman 1987](#)). The position angle (PA, hereafter) is defined as the angle between the north direction on the sky and the projected major axis of the receding half of the rings.

It is worth noting that the Eq. 1 is strictly valid only assuming that the gas is settled in a razor-thin disc and it describes only the circular motion of the gas. Deviations from pure circular orbits can include radial motions due to inflow-outflow, non-circular motions due to deviation from symmetry of the galactic potential (spiral arms, mergers, misaligned DM halo and so on; see e.g. [Schoenmakers et al. 1997](#) and [Swaters et al. 1999](#)) and small-scale perturbations due to the star formation activity (stellar winds, SNe, see e.g. [Read et al. 2016c](#)). The tilted-ring model allows us to trace the radial variation of the HI disc geometry and model the so-called ‘warp’ ([García-Ruiz et al. 2002](#); [Battaglia et al. 2006](#)). Moreover, using the ring decomposition it is possible also to obtain radial profiles from the integrated 2D maps (e.g. the HI surface density, see Sec. 4.1)

Galaxy	D (Mpc) (1)	M_V (mag) (2)	f_c (pc/arcsec) (3)	cube (4)	Beam (arcsec×arcsec) (5)	Ch. sep (km/s) (6)	M_{HI} ($10^7 M_\odot$) (7)	σ_{ch} (mJy/Beam) (8)	σ_{3T} (M_\odot/pc^2) (9)
CVnIdwA	3.6	-12.4	17.4	ro	10.9 x 10.5	1.3	4.5 ± 0.1	0.63	0.96
DDO 47	5.2	-15.5	25.2	na	16.1 x 15.3	2.6	35.3 ± 0.6	0.61	0.41
DDO 50	3.4	-16.6	16.5	na†	15.0 x 15.0	2.6	59.5 ± 1.6	0.82	0.65
DDO 52	10.3	-15.4	49.9	na	11.8 x 7.2	2.6	22.3 ± 0.1	0.46	1.20
DDO 53	3.6	-13.8	17.4	ro	6.3 x 5.7	2.6	4.7 ± 0.3	0.57	3.80
DDO 87	7.4	-15.0	35.9	na	12.8 x 11.9	2.6	20.7 ± 1.0	0.51	0.90
DDO 101	6.4	-15.0	31.0	ro	8.3 x 7.0	2.6	1.8 ± 0.2	0.50	0.90
DDO 126	4.9	-14.9	23.7	na	12.2 x 9.3	2.6	12.8 ± 0.1	0.41	1.27
DDO 133	3.5	-14.8	17.0	ro	12.4 x 10.8	2.6	10.0 ± 0.2	0.35	0.89
DDO 154	3.7	-14.2	18.0	na†	15.0x15.0	2.6	25.1 ± 0.1	0.44	0.52
DDO 168	4.3	-15.7	20.8	na	15.0 x 15.0	2.6	25.1 ± 0.1	0.47	0.54
DDO 210	0.9	-10.9	4.4	na	16.6 x 14.1	1.3	0.2 ± 0.1	0.75	0.59
DDO 216	1.1	-13.7	5.3	ro	16.2 x 15.4	1.3	0.5 ± 0.1	0.91	0.40
NGC 1569	3.4	-18.2	16.5	na†	15.0 x 15.0	2.6	17.3 ± 0.1	0.77	1.10
NGC 2366	3.4	-16.8	16.5	na†	15.0 x 15.0	2.6	64.0 ± 1.2	0.52	0.59
UGC 8508	2.6	-13.6	12.6	na†	15.0 x 15.0	1.3	1.7 ± 0.1	1.31	0.54
WLM	1.0	-14.4	4.8	na†	25.0 x 25.0	2.6	5.6 ± 0.1	2.00	0.58

Table 1. Properties of the sample of galaxies studied in this work. - (1) distance in Mpc (Hunter et al. 2012 and references therein); (2) V band absolute magnitude (Hunter et al. 2012 and references therein); (3) conversion factor from arcsec to pc; (4) weighted scheme used to produce the final datacube: "na" for natural "ro" for robust. A † means that we have further smoothed the original datacube to the beam indicated in Col. 5; (5) beam major x minor axis in arcsec of the data used for the kinematic fits (see Sec. 2.2); (6) channel separation in km/s (this value indicates also the velocity resolution of the datacube in terms of FWHM); (7) total HI mass (see Sec. 4.1.1); (8) rms noise per channel estimated with 3DB; (9) $3\text{-}\sigma$ pseudo noise in the total map (See Sec. 4.1.2). Note that DDO 87 has a slightly different distance with respect to the one reported in Hunter et al. (2012) because their value is a typo (the correct value is given in Croxall et al. 2009)

In this work we measure the kinematic and geometric properties of the galaxies in our sample with the publicly available software 3DB³ (Di Teodoro & Fraternali 2015) that is a 3D code based on the tilted-ring approach.

3.2 3DBAROLO

3DB is a 3D method that performs a tilted-ring analysis on the whole datacube. In practice, for each sampling radius it builds a ring model based on Eq. 1, then it calculates the residuals between the emission of the model and of the data pixel-by-pixel along the ring in the datacube. The final parameters of each ring are found through the minimisation of these residuals. Before the comparison between the datacube and the model, the latter is smoothed to the spatial and spectral instrumental resolution. This ensures full control of the observational effects and in particular a proper account of beam smearing that can strongly affect the derivation of the rotation velocities in the inner regions of dwarf galaxies (see e.g. Swaters 1999). Moreover 3DB fits, at the same time, the rotation velocity and the velocity dispersion instead of treating them as separate components as done in the classical 2D approach (e.g. Tamburro et al. 2009, O15).

In conclusion, 3DB fits up to 8 parameters for each ring in which the galaxy is decomposed: central coordinates, V_{sys} , i , PA, HI surface density (Σ), HI thickness (z_d), V_{rot} and the velocity dispersion (σ_v). 3DB separates the genuine HI emission from the noise by building a mask: only the pixels containing a signal above a certain threshold from the noise are taken into account. The noise estimated with 3DB for each galaxy is reported in the Col. 8 of Table 1. Further details on 3DB can be found in Di Teodoro & Fraternali (2015).

3DB works well both on high-resolution and low-resolution

data (Di Teodoro & Fraternali 2015). This allow us to use the optimal compromise between spatial resolution and sensitivity, as already discussed in Sec. 2.2. Read et al. (2016c) showed that 3DB is capable of obtaining a good estimate of the kinematic parameters of mock HI datacubes corresponding to low-mass isolated dIrrs, similar to those that we will study here.

3.2.1 Assumptions

As the focus of this work is the kinematics (V_{rot} and σ_v) of the HI discs, we can consider the other 6 variables fitted by 3DB as “nuisance parameters”. We decided to reduce the relatively high number of parameters making assumptions on the HI surface density, HI scale height, systemic velocity and the position of the galactic centre.

- *HI surface density*: we remove the surface density from the list of the free parameters by normalising the HI flux. 3DB implements two different normalisation techniques: pixel-by-pixel or azimuthally averaged. In the first case the model is normalised locally to the value of the total HI map, while in the second case the model is normalised to the azimuthally-averaged flux in each ring. For a full description of these normalisation techniques, see the 3DB reference paper (Di Teodoro & Fraternali 2015). We decided to use the local pixel-by-pixel option, because this approach is convenient to take into account the asymmetry of the HI distribution and to avoid that regions with peculiar emission (e.g. clumps or holes) affect excessively the global fit (see Lelli et al. 2012a and Lelli et al. 2012b). However, we checked that the results obtained with the two normalisations are fully compatible within the errors.
- *HI scale height*: 3DB includes the possibility to fit the HI scale height. This is something exclusive to 3D methods since there is no information about the scale height in the integrated 2D velocity field. However, the assumption of a thick disc is somewhat inconsistent with the ‘tilted-ring model’ (see Sec. 3.1): in the presence

³ <http://editeodoro.github.io/Bbarolo/>.

of a thick disc along the line of sight we are accumulating emission from different rings, hence a ring-by-ring analysis can not incorporate this effect. To be fully consistent, we should set the value of the scale height z_d to 0: however, it turns out that, assuming $z_d = 0$ the galactic models made with 3DB have a too sharp cut-off at the border of rings. Therefore, we decided to set for all the galaxies a scale height of 100 pc, constant in radius. In Sec. 7.1 we discuss in detail the effect of this assumption.

- *Systemic velocity*: we fix the systemic velocity for all rings to the value calculated as

$$V_{\text{sys}} = 0.5(V20_{\text{app}} + V20_{\text{rec}}), \quad (2)$$

where V20 is the velocity where the flux of the global HI profile is 20% of the flux peak, while ‘app’ and ‘rec’ indicate the approaching and the receding halves of the galaxy.

- *Centre of the galaxy*: We fix the position of the galactic centre for all rings to the value found with one of the methods described in Appendix A.

4 DATA ANALYSIS

4.1 HI total map

Before starting the kinematic fit procedure, for each galaxy we produce HI total maps from the datacubes. This step is needed both to have an initial rough estimate of the geometrical properties of the HI discs and to define the maximum radius to use in the kinematic fit. Moreover, we use it to extract the HI surface-density profiles using the best-fit parameters of the disc (Sec. 4.2). The HI surface-density profile is crucial to correct the estimated rotation curve for the asymmetric drift (Sec. 4.3.1).

4.1.1 HI spatial distribution

Integrating the signal of each pixel along the spectral axis one obtains a 2D total map that represents the spatial distribution (on the sky) of the HI emission. To build these maps, we masked the original datacube with the following procedure: first we smoothed the original data to obtain a low-resolution cube (between 2 and 3 times the original beam, depending on the galaxy); then, a pixel is included in the mask only if its emission is above a certain threshold (2.5 of the noise σ_{ch} of the smoothed cube) in at least three adjacent channels. Finally we sum, along the spectral axis, all the flux of the pixel included in the mask obtaining a 2D map. This method produces a very clean final map with only a small contribution from the noise.

Following Roberts (1975), we can directly relate the intensity of the radio emission (S) to the projected surface-density of the gas (Σ_{obs}) as

$$\frac{\Sigma_{\text{obs}}(x, y)}{M_{\odot}/\text{pc}^2} = 8794 \left(\frac{S(x, y)}{\text{Jy/Beam}} \right) \left(\frac{\Delta V}{\text{km/s}} \right) \left(\frac{B_{\text{maj}} B_{\text{min}}}{\text{arcsec}^2} \right)^{-1}, \quad (3)$$

where B_{maj} and B_{min} are the FWHM of the major and minor axis of the beam and ΔV is the channel separation of the datacube. We obtained the radial profile of Σ_{obs} averaging the total map along elliptical rings defined by the disc geometrical parameters (see Sec. 4.2). The errors on Σ_{obs} are calculated as the standard deviation of the averaged values⁴. The intrinsic surface density (Σ_{int}) is defined




Figure 2. Comparison of the HI mass estimated in this work and in other works (O15, blu circles; Hunter et al. (2012), red diamonds.)

as the density of the HI disc integrated along the vertical axis of the galaxy. Assuming a razor-thin HI disc, the relation between Σ_{obs} and Σ_{int} is simply

$$\Sigma_{\text{int}}(R) = \Sigma_{\text{obs}}(R) \cos i, \quad (4)$$

because the area of the projected ring is smaller than the one of the intrinsic ring by a factor $\cos i$. Finally, we can have a measure of the total mass of the HI disc summing the observed surface densities of all pixels in the total map multiplied by the physical area of the pixels

$$\frac{M_{\text{HI}}}{M_{\odot}} = 23.5 \left(\frac{\delta}{\text{arcsec}} \right)^2 \left(\frac{D}{\text{Mpc}} \right)^2 \sum_{\text{pixels}} \left(\frac{\Sigma_{\text{obs}}(x, y)}{M_{\odot}\text{pc}^{-2}} \right), \quad (5)$$

where δ is the size of the pixel and D is the distance of the galaxy. To have an estimate of the uncertainties in this measure we have built two other maps: a low-resolution map using the smoothed cube that we used to build the mask and an extremely low-resolution map smoothing again the cube to a beam of 40-60 arcsec. We applied the primary beam correction to the total HI map using the task PBCORR of GIPSY (van der Hulst et al. 1992). For each map the mass is estimated using Eq. 5 and the method described above. In particular, we take as measure of the disc mass the average and the associated error the standard deviation of the values obtained in the three maps. The final estimates of M_{HI} are listed in Table 1 (Col. 7). The intrinsic surface-density profile and the HI map contours are shown respectively in the box B (bottom right panel) and in the box C (left panel) of the summary plot of each galaxy (Sec. 5).

Figure 2 shows the comparison between the mass estimated with our method and those estimated by O15 and Hunter et al. (2012). The relative difference between our measures and those of O15 is on average 20%, with the largest discrepancies (up to 50%) found at the low-mass end; these differences are mainly due

timated errors are an underestimate of the real uncertainties (see Sicking 1997). However, the estimated errors for the surface density are already very large and we did not correct them. Moreover, the elliptical rings are usually much larger than the beam, so the correlation of the data is not expected to affect significantly the final estimate of the average and the error of the surface density.

⁴ Pixels in datacubes are correlated by the convolution with the instrumental beam occurring during the observations. As a consequence, the es-

to the different method used to estimate the total mass. The method used in [Hunter et al. \(2012\)](#) is similar to the one used in this work ([Hunter et al. 2012](#) used only robust datacubes while we used both robust and natural datacubes; see Col. 4 in Tab. 1): in this case the average relative difference is about 10%. Because of the data-reduction methods used in [Hunter et al. \(2012\)](#), the natural datacubes have systematically lower flux with respect to the robust ones (see [Hunter et al. 2012](#) for the details), therefore our values are slightly lower.

4.1.2 Noise in the total map and maximum radius

In order to define a maximum radius for the HI disc, we need to calculate the noise level in the total map. If we sum N independent channels each with noise σ_{ch} the final noise will be equal to $\sigma_{\text{ch}} \sqrt{N}$. However, using a mask, the number of summed channels is different from pixel to pixel, moreover the datacubes in our sample have been Hanning-smoothed and adjacent channels are not independent (see [Verheijen & Sancisi 2001](#)). To have a final estimate of the noise in the total map we follow the approach of [Verheijen & Sancisi \(2001\)](#) and [Lelli et al. \(2014a\)](#): we constructed a signal-to-noise map and we defined as the $3\text{-}\sigma$ pseudo level (σ_{3T}) the mean value of the pixels with a S/N between 2.75 and 3.25. We fit our kinematic model only to the portion of the datacube within the contour defined by σ_{3T} and avoiding to use rings which do not intercept, around the major axis, HI emission coming from the disc. This ensures a robust estimate of the galactic kinematics avoiding regions of the galaxy dominated by the noise and with poor information about the gas rotation. The $3\text{-}\sigma$ pseudo noise levels σ_{3T} are listed in Table 1 (Col. 9), while the maximum radii R_{max} used in the fit are reported in Table 2 (Cols. 1-2).

4.2 Datacube fit

Given the assumptions we made (Sec. 3.2.1), we are left with two geometrical (i , PA) and two kinematic (V_{rot} and σ_v) parameters.

Eq. 1 shows that the rotational velocity and the geometrical parameters are coupled: in particular they become degenerate for galaxies with rising rotation curves ([Kamphuis et al. 2015](#)) as it is the case for most of the dlrrs. As a consequence, the fitting algorithm tends to be sensitive to the initial guesses, so it is important to initialise the fit with educated guess values. We estimated i and PA using both the HI datacube and the optical data as explained in the Appendix A. The initial values for V_{rot} and σ_v have been set respectively to 30 km/s and 8 km/s for all the galaxies. 3DB allows to use an azimuthal weighting function $w(\theta)$ (see Eq. 1) to “weigh” the residuals non-uniformly across the rings (see [Di Teodoro & Fraternali 2015](#)). We decided to use $w(\theta) = \cos \theta$ to weigh most the regions around the major axis, where most of the information on the galactic rotation lies. If we fit at the same time the four parameters, we can obtain rotation curves and velocity-dispersion profiles that show unphysical discontinuities due to the scatter noise of the geometrical parameters. For this reason, we run 3DB with the option *TWO_STAGE* (see Appendix D) turned on: first a fit with four free parameters is obtained, then the geometrical parameters are regularised with a polynomial, and finally a new fit of only V_{rot} and σ_v is obtained. We chose to use the lowest polynomial order allowed by the data: in practice, we set the geometrical parameters to a constant value, unless there is a clear evidence of radial trends of i and/or of PA as for example in DDO 133 and in DDO 154. We assessed the existence of a radial trend and set the degree of the

polynomial through the visual inspection of the radial profile of i and/or PA, and also of the HI map, of the velocity field and of the datacube channels.

The result of the fit is compared by-eye with the real datacube both analysing the position-velocity diagrams (PV, hereafter) along the major axis and the minor axis, and channel by channel. Figures 3 and 4 show the comparison in the channel maps between the observations and the best-fit model found with 3DB for DDO 154 and NGC 2366: the emission of DDO 154 is reproduced quite well; the best-fit model found for NGC 2366 traces the global kinematics but it is not able to reproduce the extended feature in the North-West of the galaxy (see e.g. channel at 120 km/s). The presence of such kind of peculiarities makes the study of the dlrrs HI disc kinematics challenging and requires careful visual inspection. Whenever we notice that the model was not a good representation of the data, we restart the fit changing the initial guesses and/or varying the order of the polynomial regularisation.

Concerning the errors on the estimated parameters, it is worth spending a few words about the differences between the standard 2D approach and 3DB. In the first case the errors are a combination of (i) nominal errors coming from the tilted ring fit (typically using a Levenberg-Marquardt algorithm) and (ii) the differences between the fit of the approaching and receding sides. It is well known that errors (i) are very small and underestimate the real uncertainties: so the final errors are dominated by (ii) and the definition of this ‘asymmetric-error’ is somewhat arbitrary. 3DB uses a Monte-Carlo method to estimate the errors further exploring the space parameters around the best fit solutions (see [Di Teodoro & Fraternali 2015](#) for further details), therefore our quoted errors should be a robust and statistically significant measure of the uncertainties of the kinematic parameters.

Given that we use the errors estimated with 3DB, we did not fit separately the approaching and receding halves of the galaxy as often done in the classical 2D methods. The only exception is DDO 126: in this case the best-fit model found with 3DB using the whole galaxy does not give a good representation of the datacube because of the strong kinematic asymmetries (see Sec. 5).

4.3 The final circular velocity

Assuming that the gas is in equilibrium, the rotation velocity of the gas (V_{rot}) can be related to the galactic gravitational potential (Φ) through the radial component of the momentum equation

$$\frac{1}{\rho} \frac{\partial \rho \sigma_v^2}{\partial R} = -\frac{\partial \Phi}{\partial R} + \frac{V_{\text{rot}}^2}{R}, \quad (6)$$

where ρ is the volumetric density of the gas and σ_v is the velocity dispersion. Therefore, the observed V_{rot} is not a direct tracer of the galactic potential when the pressure support term ($\rho \sigma_v^2$) due to the random gas motions is non-negligible.

4.3.1 Asymmetric-drift correction

Eq. 6 can be re-written as

$$V_c^2 - V_{\text{rot}}^2 = -\frac{R}{\rho} \frac{\partial \rho \sigma_v^2}{\partial R} = -R \sigma_v^2 \frac{\partial \ln(\rho \sigma_v^2)}{\partial R} = V_A^2, \quad (7)$$

where $V_c = \sqrt{R \frac{\partial \Phi}{\partial R}}$ is the circular velocity and V_A is the asymmetric drift, which becomes increasingly important at large radii and is heavily dependent on the value of the velocity dispersion.




Figure 3. Channel maps showing the HI emission for DDO 154: data (top panels, blue contours) and the best-fit model found with 3DB (bottom panels, red contours). The velocity of each channel is reported above the panels. The minimum contour is $3 \sigma_{\text{ch}}$ (Col. 8 in Tab. 1) and adjacent contours differ by a factor of 2 in emissivity; the grey contours in the the data channels indicate the emission at $-3 \sigma_{\text{ch}}$. The size of the field is about $14' \times 14'$.




Figure 4. Same as Fig. 3 but for NGC 2366. The size of the field is about $20' \times 20'$.

When the rotation velocity is much higher than the velocity dispersion the asymmetric-drift correction is negligible. This is the case in spiral galaxies where usually $V_{\text{rot}}/\sigma_v > 10$ (e.g. de Blok et al. 2008). Instead, for several dIrrs the rotation velocity is comparable to the velocity dispersion, making the asymmetric-drift correction indispensable for an unbiased estimate of the galactic potential.

The volumetric density in the equatorial plane ($z=0$) is proportional to the ratio of the intrinsic surface density and the vertical scale height (z_d), therefore the asymmetric drift V_A^2 is given by

$$V_A^2 = -R\sigma_v^2 \frac{\partial \ln(\sigma_v^2 \Sigma_{\text{int}} z_d^{-1})}{\partial R}. \quad (8)$$

We can ignore the term with the radial derivative of z_d assuming that the thickness of the gaseous layer is independent of radius (but see Sec. 7.1). Furthermore, assuming that the HI disc is thin, the ratio between the intrinsic and the observed surface density is just the cosine of i (Eq. 4). Under these assumptions, from Eq. 8 we derive the classical formulation of the asymmetric-drift correction (see e.g. O15):

$$V_A^2 = -R\sigma_v^2 \frac{\partial \ln(\sigma_v^2 \Sigma_{\text{obs}} \cos i)}{\partial R}. \quad (9)$$

Except for DDO 168, NGC 1569 and UGC 8508, all the analysed galaxies have a constant i and the cosine in Eq. 9 can also be ignored.

4.3.2 Application to real data

Fluctuations of the observed surface density and of the measured velocity dispersion at similar radii can have dramatic effects on the numerical calculation of the radial derivative in Eq. 9. As a consequence, the final asymmetric-drift correction can be very scattered causing abrupt variations in the final estimate of the circular velocity. For this reason, we decided to use functional forms to describe both the velocity dispersion and the argument of the logarithm in Eq. 9. The velocity dispersion is regularised with a polynomial $\sigma_p(R, n_p)$ with degree n_p lower than 3. If there is not a clear radial trend, we consider a fixed velocity dispersion taking the median of $\sigma_v(R)$. The radial variation of the velocity dispersion is usually small, therefore the radial trend of $\Sigma_{\text{int}}\sigma_v^2$ is dominated by the behaviour of the surface density: this falls off exponentially at large radii, while in the centre it is almost constant or it shows an inner depression. In analogy with Bureau & Carignan (2002) we chose to fit $\Sigma_{\text{int}}\sigma_v^2$ with the function

$$f(R) = f_0 \left(\frac{R_c}{\text{arcsec}} + 1 \right) \left(\frac{R_c}{\text{arcsec}} + e^{\frac{R}{R_d}} \right)^{-1}, \quad (10)$$

where f_0 is a normalisation coefficient, and R_c and R_d are characteristic radii. The function f is characterised by an exponential decline at large radii and by an inner core almost equal to f_0 . Read et al. (2016c) showed that the Eq. 10 is a good compromise between a pure exponential that overestimates the asymmetric-drift correction in the inner radii, and a functional form with an inner depression that can produce unphysical negative values of V_{rot}^2 . Combining Eq. 10 and Eq. 9 we can write the asymmetric-drift correction as

$$V_A^2 = R \frac{\sigma_p^2(R, n_p) e^{\frac{R}{R_d}}}{R_d} \left(\frac{R_c}{\text{arcsec}} + e^{\frac{R}{R_d}} \right)^{-1}. \quad (11)$$

4.3.3 Error estimates

We can calculate the final errors on V_c by applying the propagation of errors to Eq. 7 to obtain

$$\delta_c = \frac{\sqrt{V_{\text{rot}}^2 \delta_{\text{rot}}^2 + V_A^2 \delta_A^2}}{V_c}, \quad (12)$$

where δ_{rot} is the error found with 3DB and δ_A is the uncertainty associated with the estimated values of the asymmetric-drift term. Usually δ_A is simply ignored (e.g. Bureau & Carignan 2002; O15), thus the uncertainties on the final corrected rotation curve are equal to δ_{rot} . This is a reasonable assumption if the rotational terms in Eq. 12 are dominant, but for some galaxies in our sample (e.g. DDO 210) the final circular rotation curve is heavily dependent on the asymmetric-drift correction. In these cases it is important that δ_c includes the uncertainties introduced by the operations described in Sec. 4.3.2. We decided to estimate δ_c with a Monte-Carlo approach:

- (i) First we make N realisations of the radial profile of both the velocity dispersion and of the surface density. For each sampling radius R the values of a single realisation $\sigma_v^i(R)$ or $\Sigma_{\text{int}}^i(R)$ are extracted randomly from a normal distribution with the centre and the dispersion taken respectively from the values and errors of the parent populations.
- (ii) For each of the N realisations we apply the method described in Sec. 4.3.2 to obtain the asymmetric-drift correction at each sampling radius $V_A^i(R)$.
- (iii) The final asymmetric-drift correction is calculated as $V_A(R) = \text{median}_i(V_A^i(R))$, where each V_A^i is obtained with the Eq. 11. The associated errors as $\delta_A(R) = K \times \text{MAD}(V_A^i(R))$ where the MAD is the median absolute deviation around the median. The factor K links the MAD with the standard deviation of the sample ($K \approx 1.48$ for a normal distribution). We chose to use the median and the MAD because they are less biased by the presence of outliers with respect to the mean and the standard deviation.

We found that $N = 1000$ is enough to obtain a good description of the error introduced by the asymmetric-drift correction.

4.3.4 Final notes

The method described in the above sections has been built to make the asymmetric-drift correction terms as smooth as possible. This ensures that our final rotation curves are not affected by non-physical noise related to the derivative term in Eq. 9. The drawback of this approach is that intrinsic scatter on V_A is hidden and the final errors δ_A could be several times larger than the point-to-point scatter. However, we are confident that the quoted values of δ_A truly trace the degree of uncertainties introduced by the asymmetric-drift correction at different radii. In galaxies where the final circular velocity is totally dependent on the asymmetric-drift correction (e.g. DDO 210, see Sec. 5 and Fig. 16) the scatter in V_{rot} could be smoothed out, but in these cases also the final errors are dominated by the errors on the asymmetric-drift and δ_c is still a good representation of the global uncertainties.

In some cases the velocity dispersion found with 3DB is not well constrained, so there are galaxies where at some radii the σ_v is discrepant or peculiar (e.g. very small error) with respect to the global trend. In general, the presence of a single ‘rogue’ σ_v (e.g. DDO 87, Fig. 10 or DDO 133, Fig. 13) does not have a significant influence on the estimate of the V_A and on the median of σ_v (see Tab. 2). However, if the discrepant σ_v is in a peculiar position (e.g. the last radius in DDO 210, Fig. 16) or the region with ‘rogues’ is

extended by more than one ring (e.g. DDO 216, empty circles in Fig. 17), the correction for the asymmetric drift and the estimate of the circular velocities are biased. In these cases we exclude the radii with discrepant σ_v both from the calculation of the asymmetric-drift correction and from the calculation of the median.

5 RESULTS

For each analysed galaxy we produced a summary plot of its properties (Figs. 5-21). Each plot is divided in four boxes: box A shows the rotation curve, the circular-velocity profile and the correction for the asymmetric drift; box B shows the velocity dispersion, the intrinsic surface density, i and PA obtained with 3DB; box C shows the total HI map and the velocity field; box D shows the PV diagrams along the major and the minor axis for both model and data. The physical radii have been calculated assuming the distance in Tab. 1 (Col. 1). A detailed description of the plot layout can be found in Appendix B. For reference, Tab. 2 lists the assumptions and the initial guesses used in 3DB (Cols. 1-7) and a summary of the results of the datacube fit (Cols. 8-11). Here below we report notes on individual galaxies. In the following notes i_{ini} and PA_{ini} indicate the initial guesses for i and PA, respectively (see Cols. 6-7 in Tab. 2).

- **CVnIdwA.** The HI morphology of CVnIdwA (Fig. 5) looks regular, but the orientation of the HI iso-density contours is quite different with respect to the direction of the velocity gradient. The emission of the stellar disc is very patchy, so we estimated the geometrical parameters of the HI disc using 3DB (see Appendix A).

- **DDO 47.** The HI disc of DDO 47 (Fig. 6) is nearly face-on, while the stellar disc looks highly inclined at 64° (Hunter & Elmegreen 2006). An inclination of 64° is not compatible with the HI contours of the total map and the stellar disc probably hosts a bar-like structure (Georgiev et al. 1997). Therefore we decided to estimate i_{ini} and PA_{ini} by fitting ellipses on the contours of the total HI map (see Appendix A). The discrepancy between the galactic centre estimated with the HI contours and the optical centre is very small (0.7 arcsec in RA and 2 arcsec in DEC), so we decided to use the HI centre. The best-fit i (see Tab. 2) is consistent with previous works (35° in Gentile et al. 2005, 30° in de Blok & Bosma 2002b and Stil & Israel 2002b) and seems a reasonable upper limit for this galaxy (see ellipses in Box C in Fig. 6). The final rotation curve shows a flattening at $R < 120''$ that could be the sign of the presence of strong non-circular motion. Moreover in the approaching half of the galaxy there is a large hole that can be clearly seen both on the total map and on the PV along the major axis: the presence of such hole could further bias the estimate of the gas rotation. For these reasons, we retain the velocities for $R < 120''$ not completely reliable (empty circles in Box A in Fig. 6).

- **DDO 50.** The gaseous disc of DDO 50 (Fig. 7) is quite peculiar: it is 'drilled' by medium-large HI holes ranging from 100 pc to 1.7 kpc (Puche et al. 1992) and it also shows clumpy regions with high density. Therefore, the HI surface density evaluated across the rings could be very discontinuous with very high peaks and/or regions without emission. As a consequence, the quoted errors on the intrinsic surface density profile (see Sec. 4.1.1) are large and they are probably over-estimating the real statistical uncertainties, especially in the inner disc ($R < 3$ kpc). Despite this peculiarity, the large scale kinematics is quite regular and typical of a flat rotation curve. The estimates of the i and PA are difficult given that this galaxy is nearly face-on. The value of i estimated with the isophotal fitting of the optical disc (47°) is too high to be compatible with

the flattening of the HI contours which favour an $i < 40^\circ$. The best agreement between the datacube and the model has been found using initial values of 30° for i_{ini} and 180° for PA_{ini} . The centre was set using the coordinates of the centre of the stellar disc (Hunter & Elmegreen 2006).

- **DDO 52.** The best-fit i of DDO 52 (Fig. 8) tends to be more edge-on at the end of the disc. However, the analysis of the channels and of the PVs indicates that the model with a constant i gives a better representation of the data.

- **DDO 53.** The stellar disc and the HI disc of DDO 53 (Fig. 9) are misaligned and the galaxy is nearly face-on. As a consequence, i and PA are very difficult to constrain: we tried different values in the range 30° - 50° for i_{ini} and 100° - 140° for PA_{ini} . The best match with the data has been found using $i_{\text{ini}} = 35^\circ$ and $\text{PA}_{\text{ini}} = 120^\circ$, but the rotational and the circular velocities should be taken with caution. In the northern part of the galaxy there is some extra emission possibly connected with an inflow/outflow: this region is clearly visible in the PV along the minor axis around -50 arcsec at a velocity of about 5 km/s.

- **DDO 87.** The morphology of DDO 87 (Fig. 10) is clearly irregular in the inner part, but the outer disc looks more regular. We decided to set the initial guesses for the PA using 3DB (see Appendix A). We tried different initial guesses for i : the best representation of the data has been obtained with $i_{\text{ini}} = 40^\circ$ and $\text{PA}_{\text{ini}} = 240^\circ$, approximately 15° lower than the orientation of the optical disc (Hunter & Elmegreen 2006).

- **DDO 101.** The HI disc of DDO 101 (Fig. 11) is extended only slightly beyond the optical disc and the HI emission is almost constant with some high density structures around 1.0 kpc. Notice that the estimates of the distance for this galaxy are very uncertain (ranging from 5 to 16 Mpc) since they all rely on a poor distance estimator (Tully-Fisher relation, e.g. Karachentsev et al. 2013).

- **DDO 126.** DDO 126 (Fig. 12) shows kinematic asymmetries, so we separately run 3DB also on the approaching and the receding halves of the galaxy. Beyond 1.5 kpc the fit on the whole galaxy gives a good representation of the datacube and the errors found with 3DB are larger than (or comparable to) the differences due to the kinematic asymmetries (black circles in Box A in Fig. 12). The inner regions are less regular and the best model has been found taking the mean between the approaching and receding rotation curves (empty circles in Box A in Fig. 12), while the errors have been calculated as half the difference between the two values following the recipe of Swaters (1999).

- **DDO 133.** The HI disc of DDO 133 (Fig. 13) has a regular kinematics, although there is evidence of non-circular motions, especially in the region of the stellar bar (Hunter et al. 2011). The final PA found with 3DB is about 20° in the inner part of the disc ($R < 0.5$ kpc) and it becomes almost zero in the outer disc. We decided to take the radial trend of the PA into account with a fourth order polynomial.

- **DDO 154.** The HI morphology and the kinematics of DDO 154 (Fig. 14) is quite regular. However, the contours of the HI map clearly show a radial trend of the PA, as confirmed by the values found with 3DB. In order to obtain a good description of the radial trend of the PA we used a fourth order polynomial. However, the best-fit polynomial shows a very steep gradient (about 25°) in a very small region (less than 1 kpc) that is not justified by the data (red dashed line in Box B in Fig. 14). The points located in this region are compatible with a constant PA, so we decided to fix it to the mean values of the first three points (see Box B in Fig. 14). As in O15, we do not confirm the almost Keplerian fall-off of the

Galaxy	R_{\max} (arcsec)	R_{\max} (kpc)	ΔR (pc)	centre		V_{sys} (km/s)	i_{ini} ($^{\circ}$)	PA_{ini} ($^{\circ}$)	V_0 (km/s)	$\langle \sigma_v \rangle$ (km/s)	$\langle i \rangle$ ($^{\circ}$)	$\langle PA \rangle$ ($^{\circ}$)
	(1)	(2)	(3)	RA (4a)	DEC (4b)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
CVnIwA	90	1.6	170	12 38 40.2 ^B	32 45 52 ^B	307.9 ^B	41 ^B	55 ^B	21.5 \pm 3.9	7.7 \pm 0.5	49.2 \pm 10.9	60.6 \pm 18.5
DDO 47	210	5.3	380	7 41 54.6 ^H	16 48 10 ^H	272.8 ^B	35 ^H	310 ^H	62.6 \pm 5.2 [†]	7.9 \pm 0.2	37.4 \pm 1.7 [†]	317.4 \pm 7.3
DDO 50	390	6.4	190	8 19 08.7 ^S	70 43 25 ^S	156.7 ^B	30 ^T	180 ^T	38.7 \pm 10.1 [†]	9.0 \pm 1.5	33.1 \pm 4.6 [†]	174.9 \pm 5.9
DDO 52	108	5.4	450	8 28 28.5 ^S	41 51 21 ^S	396.2 ^B	51 ^S	5 ^S	51.1 \pm 6.3	8.0 \pm 0.8	55.1 \pm 2.9	6.5 \pm 2.0
DDO 53	60	1.1	110	8 34 08.0 ^S	66 10 37 ^S	20.4 ^B	35 ^T	120 ^T	20.3 \pm 6.6 [†]	7.9 \pm 0.1	37.0 \pm 2.0 [†]	123.4 \pm 3.8
DDO 87	144	5.2	430	10 49 34.7 ^S	65 31 46 ^S	338.7 ^B	40 ^T	240 ^B	50.3 \pm 9.1	6.3 \pm 2.4	42.7 \pm 7.3	238.6 \pm 4.7
DDO 101	60	1.9	190	11 55 39.4 ^S	31 31 8 ^S	586.6 ^B	49 ^S	290 ^S	59.2 \pm 3.6	4.8 \pm 1.7	52.4 \pm 1.7	287.4 \pm 3.6
DDO 126	140	3.3	240	12 27 06.3 ^H	37 08 23 ^H	214.3 ^B	63 ^B	138 ^B	38.6 \pm 3.1	9.1 \pm 1.6	62.2 \pm 2.9	140.7 \pm 3.5
DDO 133	165	2.8	190	12 32 55.4 ^S	31 32 14 ^S	331.3 ^B	38 ^B	20 ^B	47.2 \pm 5.1 [†]	8.1 \pm 0.7	38.9 \pm 3.7 [†]	-3.8 \pm 8.9
DDO 154	390	7.0	270	12 54 06.2 ^S	27 09 02 ^S	375.2 ^B	65 ^S	224 ^S	47.1 \pm 5.1	8.5 \pm 0.9	67.9 \pm 1.1	226.1 \pm 2.6
DDO 168	225	4.7	310	13 14 27.9 ^B	45 55 24 ^B	191.9 ^B	60 ^T	300 – 270 ^T	56.2 \pm 6.9	8.8 \pm 1.3	62.0 \pm 2.6	272.7 \pm 4.2
DDO 210	100	0.4	40	29 46 52.0 ^S	-12 59 51 ^S	-140.0 ^B	60 ^T	65 ^T	16.4 \pm 9.5	6.2 \pm 0.6	63.2 \pm 3.2	77.3 \pm 15.2
DDO 216	195	1.0	80	23 28 32.1 ^H	14 44 50 ^H	-188.0 ^T	65 ^H	130 ^H	13.6 \pm 5.5	5.6 \pm 0.5	70.0 \pm 5.0	130.4 \pm 9.0
NGC 1569	150	2.5	250	4 30 49.8 ^S	64 50 51 ^S	-75.6 ^B	61 ^S	122 ^S	55.6 \pm 22.4	21.0 \pm 4.0	67.0 \pm 5.6	114.6 \pm 4.0
NGC 2366	384	6.3	260	7 28 48.8 ^S	69 12 22 ^S	100.8 ^B	65 ^B	35 ^B	57.7 \pm 5.4	12.6 \pm 1.8	65.1 \pm 4.2	39.8 \pm 2.8
UGC 8508	110	1.4	130	13 39 44.9 ^S	54 54 29 ^S	59.9 ^B	65 ^H	120 ^H	33.8 \pm 6.4	9.1 \pm 1.8	67.6 \pm 5.3	123.2 \pm 1.7
WLM	600	2.9	120	0 01 59.2 ^S	-15 27 41 ^S	-124.0 ^B	70 ^S	178 ^S	38.7 \pm 3.4	7.7 \pm 0.8	74.0 \pm 2.3	174.0 \pm 3.1

Table 2. Fit assumptions: - (1) outermost radius in arcsec (see Sec. 4.1.2); (2) outermost radius in kpc (see Sec. 4.1.2); (3) sampling radius used in the datacube fitting; (4a)(4b) assumed galactic centre in RA (hh mm ss.s) and DEC (dd mm ss) coordinate system (J2000); (5) assumed systemic velocity (see Sec. 3.2.1); (6) initial guesses for i ; (7) initial guesses for PA. ^S indicates that the values is estimated from the stellar disc (Hunter & Elmegreen 2006), ^B using 3DB, ^H using the fit of the contours of the HI total map (see Appendix A for details). ^T indicates peculiar cases (see the individual galactic description for details). **Results:** (8) mean circular velocity of the outer disc (see Sec. 6); (9) median of the best-fit velocity dispersion found with 3DB; (10) median of the best-fit PA found with 3DB; (11) median of the best-fit i found with 3DB. [†] indicates nearly face-on galaxies in which the quoted errors on $\langle i \rangle$ and V_0 could underestimate the real uncertainties.

rotation curve claimed by Carignan & Purton (1998) beyond 5 kpc. The channel maps are shown in Fig. 3

- **DDO 168.** The velocity gradient of DDO 168 (Fig. 15) is misaligned with respect to both the HI and the stellar disc. Moreover, the presence of a prominent bar visible both in the stellar disc (Hunter & Elmegreen 2006) and in the inner part of the HI disc makes the initial estimate of the geometrical parameters very uncertain. We obtained a first estimate of the PA using the 2D tilted-ring fitting of the velocity field shown in Fig. 15: the resultant PA decreases from 300° to about 270°. We used these values as initial guesses for all sampling radii in 3DB (see Appendix A). The centre was set to the value found with 3DB which roughly corresponds to the optical centre. We tried different values for i_{ini} between 40° – 70°: the best reproduction of the datacube has been obtained with $i_{\text{ini}} = 60^\circ$. The final results still show a variation of PA that we regularised with a second order polynomial. The outer disc of DDO 168 is quite irregular. There is extra emission at velocities close to V_{sys} possibly related to inflow/outflow, while the distortions of the iso-velocity contours (Box C. in Fig. 15) could be due to the presence of an outer warp.

- **DDO 210.** DDO 210 (also known as Aquarius dIrr, Fig. 16) is the least massive galaxy in our sample and it is classified as a transitional dwarf galaxy (McConnachie 2012 and reference therein). The HI map is quite peculiar with isodensity contours that are not elliptical. As a consequence the estimate of the galactic centre using the HI data is very uncertain and we decided to set it to the optical value (Hunter & Elmegreen 2006). The kinematics is dominated by the velocity dispersion, however a weak velocity gradient is visible. The velocity gradient looks misaligned with both the stellar and the HI disc, so we set the initial values of i and PA (60° and 65°, respectively) using a by-eye inspection of the velocity field. This procedure is arbitrary, but it is important to note that the final circular-velocity curve and the related errors are independent of our procedure since they are totally dominated by the asymmetric-drift correction (see Sec. 4.3.1 and Sec. 4.3.4). The V_{rot} found with 3DB

in the first two rings (10 and 20 arcsec) is not well-constrained, so we preferred to exclude them. Notice that along the minor axis there is an extended region with HI emission apparently not connected with the rotating disc. As in the case of DDO 53 this emission could trace an inflow/outflow.

- **DDO 216.** DDO 216 (also known as Pegasus dIrr, Fig. 17) is defined as a transitional dwarf galaxy (Cole et al. 1999). The galaxy shows a velocity gradient aligned with the HI and the optical disc, but it could be entirely due to a single ‘cloud’ at a discrepant velocity in the approaching side of the galaxy (Stil & Israel 2002a). For the purpose of our work, we assumed the gradient genuine and caused entirely by the gas rotation. The analysis of the alternative scenario can be found in Appendix C. Given the kinematic peculiarities (see Appendix C), we did not use 3DB to estimate the V_{sys} (Sec. 3.2.1). We tried different values between -186 and -190 km/s and we decided to use $V_{\text{sys}} = -188$ km/s because this value minimises the kinematic asymmetries between the receding and the approaching halves of the galaxy. We set the centre, i_{ini} (65°) and PA_{ini} (130°) using the values obtained from the elliptical fit of the outermost HI contours ($R > 600$ pc). The best-fit PA decreases from about 140° to about 125°.

- **NGC 1569** NGC 1569 (Fig. 18) is a starburst galaxy with a very disturbed HI kinematics and morphology (Stil & Israel 2002b; Johnson et al. 2012; Lelli et al. 2014b). We found that the best-fit model shows a slight increase of i and a slight decrease of PA as a function of radius. The best-fit model is a good representation of the large scale structure and kinematics of the HI disc, but it fails to reproduce the small scale local features. The ISM of this galaxy is highly turbulent: the velocity dispersion found with 3DB is about 20 km/s and the asymmetric-drift correction dominates at all radii. For this reason the kinematic data reported here should be used with caution especially at the inner radii (empty circles in Box A in Fig. 18) where no significant rotation of the gas is observed (see also Lelli et al. 2014b).

- **NGC 2366.** The HI disc of NGC 2366 (Fig. 19) is quite regular, although it shows some peculiar features. The HI emission on the channel maps (Fig. 4) indicates the presence of two ridges located in the North-West and South-East (less prominent) of the disc running parallel to the major axis. The ridges show different kinematics with respect to the disc (see Oh et al. 2008) and their origin is not clear (see Hunter et al. 2001 for a detailed discussion). We checked that 3DB was not affected by the presence of this feature. From the PV along the major axis (Panel D in Fig. 19) it is clear that the 3DB model does not reproduce some emission close to the systemic velocity, especially in the receding side where the gas is seen also at ‘forbidden’ velocities below V_{sys} . As already stated by Lelli et al. (2014b) this is probably due to the presence of some extraplanar gas that is rotating at lower velocity with respect to the gas in the disc (see e.g. Fraternali et al. 2002). As in Lelli et al. (2014b) and in O15 we do not find that the rotation curve declines beyond 5 kpc, as instead claimed by both Hunter et al. (2001) and van Eymeren et al. (2009a).

- **UGC 8508.** The HI and the stellar disc of UGC 8508 (Fig. 20) are aligned but the analysis of the HI map favours an i slightly higher than the value obtained from the stellar disc (Hunter & Elmegreen 2006). We found that the datacube is better reproduced with a linearly increasing i . Notice that in the inner part of the galaxy ($R < 0.6$ kpc) the kinematics is very peculiar as it is visible by the S-shaped iso-velocity contours (right-panel C in Fig. 20). This kind of distortions can be related to an abrupt variation of the PA and/or to the presence of radial motions (Fraternali et al. 2001) as well as to a deviation from axisymmetry of the galactic potential (Swaters et al. 1999). We tested the hypothesis of a radially varying PA and the presence of non-zero radial velocities (V_{rad}) performing a 2D analysis of the velocity field with ROTCUR (Begeman 1987). We found that the combination of the two effects can partially explain the distortions of the velocity field, but their magnitude is too large to be physically plausible. Fortunately, the final rotation curves obtained including radial motions and/or the varying PA are compatibles with the results we found with 3DB, though the inner points (empty circles in Box A in Fig. 20) should be treated with caution.

- **WLM.** The HI and the optical discs of WLM (Fig. 21) are well aligned, but the best-fit i looks slightly too edge-on with respect to the HI contours (see ellipses in Fig. 21). The excess of the emission around the minor axis could be partially due to the thickness of the gaseous layer (Sec. 7.1, see also Leaman et al. 2012). Further details on the analysis of WLM can be found in Read et al. (2016c).

6 APPLICATION: TEST OF THE BARYONIC TULLY-FISHER RELATION

The baryonic Tully-Fisher relation (BTFR) links a characteristic circular velocity (V) of a galaxy with its total baryonic mass (M_{bar}). The relation, in the logarithmic form

$$\log\left(\frac{M_{\text{bar}}}{M_{\odot}}\right) = s \log\left(\frac{V}{\text{km s}^{-1}}\right) + A \quad (13)$$

is very tight and extends over 6 decades in M_{bar} (McGaugh 2012; Lelli et al. 2016b). The existence of this relation represents a fundamental benchmark for cosmological models and for galaxy formation theories (McGaugh et al. 2000; Di Cintio & Lelli 2016; Brook et al. 2016b). In this context it is very important to extend the study of the BTFR down to extremely low-mass dwarf galaxies (e.g. Begum et al. 2008). In this section we present the BTFR for the galaxies in our sample (Fig. 22) and then we compare it with

the results of Lelli et al. (2016b) (Fig. 23). The derivation of the baryonic mass and of the characteristic velocity is described below.

• **Rotation velocity.** The rotation velocities used in the Tully-Fisher relation (TFR) are W_{20} and V_{flat} : W_{20} is the velocity width of the global HI line profiles, while V_{flat} is the value of the flat portion of the rotation curve. Verheijen (2001) found that V_{flat} minimises the scatter in the TFR, so it is the best way to study the BTFR using high resolution HI data (see also Brook et al. 2016b). However, several rotation curves of dIrrs do not reach the flat part (e.g. DDO 53, Fig. 9) or the flattening is entirely due to the asymmetric-drift correction (e.g. NGC 1569, Fig. 18). As indicator of the rotation velocity, we therefore used the velocity of the outer disc (V_o) defined as the mean circular velocity of the last three fitted rings (see Sec. 4.2). V_o is a measure of V_{flat} for galaxies in which the flat part of the rotation curve is observed. For galaxies with a rising rotation curve V_o is an estimate of the maximum circular velocity within the considered radial range. Galaxies with rising rotation curves are indicated with empty markers in Figs. 22 and 23. The error on V_o is conservatively assumed as the maximum error between the three values used to calculate V_o . In the five galaxies with a low i (squares in Figs. 22 and 23) the uncertainties on i and on the velocities can be largely underestimated (see Sec. 5 and Tab. 2), so we present these galaxies with a bar that indicates the interval between a minimum (assuming $i = 40^\circ$) and a maximum (assuming $i = 20^\circ$) value of V_o (see Eq. 1).

• **Baryonic mass.** The total baryonic mass of the galaxies has been calculated as

$$M_{\text{bar}} = M_* + 1.33M_{\text{HI}}, \quad (14)$$

where M_* is the stellar mass, M_{HI} is the mass of the atomic hydrogen and the factor 1.33 takes into account the presence of Helium (Begum et al. 2008; Lelli et al. 2016a). The molecular gas is likely irrelevant in the mass budget of dwarf galaxies (Taylor et al. 1998). The mass of atomic gas is measured from the HI datacubes (Sec. 4.1.1 and Tab. 1), while the stellar masses are from Walter & Brinks (2001) for DDO 47 and from Zhang et al. (2012) for all the other galaxies. The estimate of the mass is proportional to the square of the galactic distance, therefore errors on the distance add further uncertainties on the final estimate of the baryonic mass. Unfortunately, the works we used to take the stellar masses (Walter & Brinks 2001; Zhang et al. 2012) and the distances (Hunter et al. 2012) do not report the errors on their measures, so we assumed a conservative error of the 30% for M_{bar} . We also performed a deeper analysis for the distance uncertainties. The relative difference between the masses estimated assuming two different distances is

$$\delta_D = \frac{M(D_1) - M(D_2)}{M(D_1)} = \frac{D_1^2 - D_2^2}{D_1^2}. \quad (15)$$

For each galaxy in our sample we choose the best distance estimator⁵ available on NED (NASA/IPAC Extragalactic Database) and we considered the minimum ($D_{\text{NED}}^{\text{min}}$) and the maximum ($D_{\text{NED}}^{\text{max}}$) estimate of the distance, then using Eq. 15 we calculated

$$\delta_D^{\text{min/max}} = 1 - \left(\frac{\min(D, D_{\text{NED}}^{\text{min/max}})}{\max(D, D_{\text{NED}}^{\text{min/max}})} \right)^2, \quad (16)$$

⁵ The scale of distance estimators is, from the best to the worst: Cepheids, RGB-Tip, CMD, Brightest-Stars, Tully-Fisher relation.




Figure 5. See caption in Appendix B. Notes. **C (left-hand panel):** Contours at $2^n \sigma_{3T}$ and $\sigma_{3T} = 0.96 M_{\odot} \text{pc}^{-2}$ (thick contour), stellar map in SDSS r band (Baillard et al. 2011); **C (right-hand panel):** Contours at $V_{\text{sys}} \pm \Delta V$ where $\Delta V = 6.0$ km/s and $V_{\text{sys}} = 307.9$ km/s (thick contour). **D:** Contours at $(2 + 4n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 0.63 \text{ mJy } \text{bm}^{-1}$, the grey contours are at $-2\sigma_{\text{ch}}$.




Figure 6. See caption in Appendix B. Notes. **A:** The empty circles indicate the region in which the estimate of the rotation curve can be biased by small HI regions at anomalous velocity along the major axis (see the text for further details); **C (left-hand panel):** Contours at $2^{\text{th}}\sigma_{3T}$ and $\sigma_{3T} = 0.41 M_{\odot}\text{pc}^{-2}$ (thick contour), stellar map in SDSS r band (Baillard et al. 2011); **C (right-hand panel):** Contours at $V_{\text{sys}} \pm \Delta V$ where $\Delta V = 20.0$ km/s and $V_{\text{sys}} = 272.8$ km/s (thick contour). **D:** Contours at $(2 + 6n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 0.61 \text{ mJy } \text{bm}^{-1}$, the grey contours are at $-2\sigma_{\text{ch}}$.




Figure 7. See caption in Appendix B. Notes. **C (left-hand panel):** Contours at $3^n \sigma_{3T}$ and $\sigma_{3T} = 0.65 M_{\odot} \text{pc}^{-2}$ (thick contour), stellar map in R band from Cook et al. (2014); **C (right-hand panel):** Contours at $V_{\text{sys}} \pm \Delta V$ where $\Delta V = 12.0 \text{ km/s}$ and $V_{\text{sys}} = 156.7 \text{ km/s}$ (thick contour). **D:** Contours at $(2 + 7n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 0.82 \text{ mJy bm}^{-1}$, the grey contours are at $-2\sigma_{\text{ch}}$.




Figure 8. See caption in Appendix B. Notes. **C (left-hand panel):** Contours at $2^n \sigma_{3T}$ and $\sigma_{3T} = 1.20 M_{\odot} \text{pc}^{-2}$ (thick contour), stellar map in SDSS r band (Baillard et al. 2011); **C (right-hand panel):** Contours at $V_{\text{sys}} \pm \Delta V$ where $\Delta V = 10.0 \text{ km/s}$ and $V_{\text{sys}} = 396.2 \text{ km/s}$ (thick contour). **D:** The contours start from $(2 + 4n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 0.46 \text{ mJy bm}^{-1}$, the grey contours are at $-2\sigma_{\text{ch}}$.




Figure 9. See caption in Appendix B. Notes. **C (left-hand panel):** Contours at $2^n \sigma_{3T}$ and $\sigma_{3T} = 3.80 M_{\odot} \text{pc}^{-2}$ (thick contour), stellar map in R band from Cook et al. (2014); **C (right-hand panel):** Contours at $V_{\text{sys}} \pm \Delta V$ where $\Delta V = 8.0 \text{ km/s}$ and $V_{\text{sys}} = 20.4 \text{ km/s}$ (thick contour). **D:** Contours at $(2 + 4n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 0.57 \text{ mJy bm}^{-1}$, the grey contours are at $-2\sigma_{\text{ch}}$.




Figure 10. See caption in Appendix B. Notes. **C (left-hand panel):** Contours at $2^n \sigma_{3T}$ and $\sigma_{3T} = 0.90 M_{\odot} \text{pc}^{-2}$ (thick contour), stellar map in SDSS r band (Baillard et al. 2011); **C (right-hand panel):** Contours at $V_{\text{sys}} \pm \Delta V$ where $\Delta V = 10.0 \text{ km/s}$ and $V_{\text{sys}} = 338.7 \text{ km/s}$ (thick contour). **D:** Contours at $(2 + 4n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 0.51 \text{ mJy } \text{bm}^{-1}$, the grey contours are at $-2\sigma_{\text{ch}}$.




Figure 11. See caption in Appendix B. Notes. **C (left-hand panel):** Contours at $4^n \sigma_{3T}$ and $\sigma_{3T} = 0.90 M_{\odot} \text{pc}^{-2}$ (thick contour), stellar map in SDSS r band (Baillard et al. 2011); **C (right-hand panel):** Contours at $V_{\text{sys}} \pm \Delta V$ where $\Delta V = 20.0 \text{ km/s}$ and $V_{\text{sys}} = 586.6 \text{ km/s}$ (thick contour). **D:** Contours at $(2 + 6n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 0.50 \text{ mJy bm}^{-1}$, the grey contours are at $-2\sigma_{\text{ch}}$.




Figure 12. See caption in Appendix B. Notes. **A:** The empty circles indicates the region in which the circular velocity has been calculated as the mean between the approaching and receding rotation curves; **C (left-hand panel):** Contours at $2^{\text{th}} \sigma_{3T}$ and $\sigma_{3T} = 1.27 M_{\odot} \text{pc}^{-2}$ (thick contour), stellar map in SDSS r band (Baillard et al. 2011); **C (right-hand panel):** Contours at $V_{\text{sys}} \pm \Delta V$ where $\Delta V = 8.0 \text{ km/s}$ and $V_{\text{sys}} = 214.3 \text{ km/s}$ (thick contour). **D:** Contours at $(2 + 6n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 0.41 \text{ mJy bm}^{-1}$, the grey contours are at $-2\sigma_{\text{ch}}$.




Figure 13. See caption in Appendix B. Notes. **C (left-hand panel):** Contours at $2^n\sigma_{3T}$ and $\sigma_{3T} = 0.89 M_{\odot}\text{pc}^{-2}$ (thick contour), stellar map in R band from Cook et al. (2014); **C (right-hand panel):** Contours at $V_{\text{sys}} \pm \Delta V$ where $\Delta V = 8.0$ km/s and $V_{\text{sys}} = 331.3$ km/s (thick contour). **D:** Contours at $(2 + 5n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 0.35$ mJy bm^{-1} , the grey contours are at $-2\sigma_{\text{ch}}$.




Figure 14. See caption in Appendix B. Notes. **B (upper-left panel):** The red dashed line shows the best-fit polynomial to the data, the red continuous line shows the regularisation of the radial profile used for this galaxy (see text for details); **C (left-hand panel):** Contours at $2^n \sigma_{3T}$ and $\sigma_{3T} = 0.52 \text{ M}_\odot \text{pc}^{-2}$ (thick contour), stellar map in R band from [Cook et al. \(2014\)](#); **C (right-hand panel):** Contours at $V_{\text{sys}} \pm \Delta V$ where $\Delta V = 10.0 \text{ km/s}$ and $V_{\text{sys}} = 375.2 \text{ km/s}$ (thick contour). **D:** Contours at $(2 + 8n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 0.44 \text{ mJy bm}^{-1}$, the grey contours are at $-2\sigma_{\text{ch}}$.




Figure 15. See caption in Appendix B. Notes. **C (left-hand panel):** Contours at $3^n \sigma_{3T}$ and $\sigma_{3T} = 0.54 M_{\odot} \text{pc}^{-2}$ (thick contour), stellar map in r band from (Baillard et al. 2011); **C (right-hand panel):** Contours at $V_{\text{sys}} \pm \Delta V$ where $\Delta V = 15.0 \text{ km/s}$ and $V_{\text{sys}} = 191.9 \text{ km/s}$ (thick contour). **D:** Contours at $(2 + 15n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 0.47 \text{ mJy } \text{bm}^{-1}$, the grey contours are at $-2\sigma_{\text{ch}}$.




Figure 16. See caption in Appendix B. Notes. **C (left-hand panel):** Contours at $2^3\sigma_{3T}$ and $\sigma_{3T} = 0.59 M_{\odot}\text{pc}^{-2}$ (thick contour), stellar map in R band from Cook et al. (2014); **C (right-hand panel):** Contours at $V_{\text{sys}} \pm \Delta V$ where $\Delta V = 2.5$ km/s and $V_{\text{sys}} = -140.0$ km/s (thick contour). **D:** Contours at $(2 + 4n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 0.75 \text{ mJy } \text{bm}^{-1}$, the grey contours are at $-2\sigma_{\text{ch}}$.




Figure 17. See caption in Appendix B. Notes. **C (left-hand panel):** Contours at $2^n \sigma_{3T}$ and $\sigma_{3T} = 0.40 M_{\odot} \text{pc}^{-2}$ (thick contour), stellar map in SDSS r band (Knapen et al. 2014); **C (right-hand panel):** Contours at $V_{\text{sys}} \pm \Delta V$ where $\Delta V = 5.0 \text{ km/s}$ and $V_{\text{sys}} = -188.0 \text{ km/s}$ (thick contour). **D:** Contours at $(2 + 6n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 0.91 \text{ mJy } \text{bm}^{-1}$, the grey contours are at $-2\sigma_{\text{ch}}$.




Figure 18. See caption in Appendix B. Notes. **A:** The empty circles indicate the region in which there are no significant signs of gas rotation; **C (left-hand panel):** Contours at $4^n\sigma_{3T}$ and $\sigma_{3T} = 1.10 M_{\odot}\text{pc}^{-2}$ (thick contour), stellar map in J band from (Jarrett et al. 2003); **C (right-hand panel):** Contours at $V_{\text{sys}} \pm \Delta V$ where $\Delta V = 15.0$ km/s and $V_{\text{sys}} = -75.6$ km/s (thick contour). **D:** Contours at $(2 + 8n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 0.77$ mJy bm^{-1} , the grey contours are at $-2\sigma_{\text{ch}}$.




Figure 19. See caption in Appendix B. Notes. **C (left-hand panel):** Contours at $3^3\sigma_{3T}$ and $\sigma_{3T} = 0.59 \text{ M}_\odot \text{pc}^{-2}$ (thick contour), stellar map in R band from Cook et al. (2014); **C (right-hand panel):** Contours at $V_{\text{sys}} \pm \Delta V$ where $\Delta V = 10.0 \text{ km/s}$ and $V_{\text{sys}} = 100.8 \text{ km/s}$ (thick contour). **D:** Contours at $(2 + 10n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 0.52 \text{ mJy bm}^{-1}$, the grey contours are at $-2\sigma_{\text{ch}}$.




Figure 20. See caption in Appendix B. Notes. **A:** The empty circles indicate the region of the galaxy with peculiar kinematics (see the text); **C (left-hand panel):** Contours at $3^n \sigma_{3T}$ and $\sigma_{3T} = 0.54 M_{\odot} \text{pc}^{-2}$ (thick contour), stellar map in R band from Cook et al. (2014); **C (right-hand panel):** Contours at $V_{\text{sys}} \pm \Delta V$ where $\Delta V = 10.0$ km/s and $V_{\text{sys}} = 59.9$ km/s (thick contour). **D:** Contours at $(2 + 6n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 1.31 \text{ mJy bm}^{-1}$, the grey contours are at $-2\sigma_{\text{ch}}$.




Figure 21. See caption in Appendix B. Notes. **C (left-hand panel):** Contours at $2^3\sigma_{3T}$ and $\sigma_{3T} = 0.58 M_{\odot}\text{pc}^{-2}$ (thick contour), stellar map in R band from Cook et al. (2014); **C (right-hand panel):** Contours at $V_{\text{sys}} \pm \Delta V$ where $\Delta V = 20.0 \text{ km/s}$ and $V_{\text{sys}} = -124.0 \text{ km/s}$ (thick contour). **D:** Contours at $(2 + 4n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 2.00 \text{ mJy bm}^{-1}$, the grey contours are at $-2\sigma_{\text{ch}}$.




Figure 22. BTFR for the galaxies in our sample. The empty and the solid symbols indicate galaxies with rising or flat rotation curve, respectively. The squares indicate nearly face-on galaxies. The blue bars indicate $1\text{-}\sigma$ errors. The light-blue and the magenta bars indicate intervals of allowed values (see text).




Figure 23. Same data as in 22 compared with the galaxies and the BTFR from Lelli et al. (2016b).

where D is the distance assumed in this work (see Tab. 1). When δ_D is large the error on the total mass is dominated by the uncertainty on the distance. Three galaxies have δ_D larger than the 60%: DDO 47 ($\delta_D^{\max} = 62\%$), DDO 101 ($\delta_D^{\max} = 85\%$) and NGC 1569 ($\delta_D^{\min} = 68\%$). For these galaxies we do not show the $1\text{-}\sigma$ error on the baryonic mass, but a magenta bar indicating the interval of mass found assuming the distance D or the distance $D_{\text{NED}}^{\min/\max}$.

In Fig. 23 we compare our data with a recent fit to the BTFR (Lelli et al. 2016b). Interestingly our data overlap between 10^8 and $10^9 M_\odot$. In this range our data are perfectly compatible with the parameters of the BTFR estimated in Lelli et al. (2016b) ($s = 3.95$ and $A = 1.86$ in Eq. 13); we also confirm the relatively small scatter around the relation, with little increase towards lower mass galaxies, in contrast to Begum et al. (2008). This remains the case, even when including galaxies with rising rotation curves: the only two outliers are DDO 50 (a nearly face-on galaxy) and DDO 101 (for

which the uncertainty on the distance is large; Sec. 5, see also Read et al. 2016c). Below $10^8 M_\odot$ the distribution of the galaxies looks again compatible with the relation of Lelli et al. (2016b). It appears more scattered, but this could owe entirely to the large error bars for these low-mass systems (that results from the increasingly important asymmetric-drift correction). Furthermore, there are few galaxies in this mass range and all of them have rising rotation curves (with the exception of DDO 216 that has rather peculiar kinematics; see Fig. 17, Sec. 5 and Appendix C).

7 DISCUSSION

7.1 HI scale height

All the galaxies in our sample have been analysed assuming a very thin HI disc with scale height 100 pc, independent of radius (see Sec. 3.2.1). This assumption would be fully justified in the case of spirals (e.g. Brinks & Burton 1984; Olling 1996), but in the shallow potential of the dIrrs the gaseous layer can be quite thick especially in the outer regions of the disc (see e.g. O’Brien et al. 2010; Roychowdhury et al. 2010). In this Section we discuss how the presence of a thick disc could bias the results of our analysis.

In the presence of a thin disc the observed HI emission can be easily related to the intrinsic properties of the galaxy: the observed velocity is a measure of gas rotation in the equatorial plane, the observed velocity dispersion is an unbiased measure of the chaotic motion of the gas and the intrinsic profile of the HI surface density can be obtained by simply correcting the observed profile for the i of the galaxy (Eq. 4). In the presence of thick gaseous layers, the line of sight intercepts the emission coming from rings at different radial and vertical positions. As a consequence, the parameters obtained by assuming zero thickness may not be a precise measure of the kinematics of the galaxy.

2D methods as in O15 work on integrated maps (velocity fields) and they can not take into account the presence of a thick disc. 3DB is more promising since the scale height is one of the parameters needed in the datacube fitting. We performed several tests with 3DB, but we found that the fit is essentially insensitive of the HI thickness for small-medium values of the scale height ($z_d \lesssim 0.7$ kpc) and returns unacceptable results for very thick HI discs. The reason for this is that 3DB fits one single ring at the time and it can not include the emission coming from the extended vertical layers of the other rings.

We quantified the magnitude and the type of errors introduced by the assumption of thin disc as follows. We performed several tests on mock datacubes made with the task GALMOD (Sicking 1997) of the software package GIPSY (van der Hulst et al. 1992). We found that the assumption of a thin disc biases the results as follows:

- (i) the surface-density profile tends to be shallower than the real profile;
- (ii) the measured broadening of the HI line profiles is larger than the intrinsic velocity dispersion of the gas (σ_v is overestimated);
- (iii) the representative velocity estimated from the HI line profiles may not trace the gas rotation velocity at certain locations.

Obviously, these effects influence the estimate of the final rotation curve: due to (iii) the observed velocities are no longer tracing the rotation on the equatorial plane, while (i) and (ii) bias the calculation of the asymmetric-drift correction (see Sec. 4.3.1). We found that the combination of these effects causes the circular velocities,

obtained with both 2D and 3D methods assuming a thin disc, to be lower than the intrinsic velocities at smaller radii (similar to the beam smearing) and higher in the outermost disc. The magnitude of these differences depends mainly on the thickness of the HI discs, but the i of the disc and the shape of the rotation curve are also important parameters. The biases described above are negligible for nearly face-on galaxies ($i < 50^\circ$), while a rising rotation curve, typical of dIrrs, amplifies the errors.

The thickness of the HI disc has never been incorporated in the derivation of the rotational velocity. In Iorio et al. (in prep) we will present an original method to estimate in a self-consistent way the intrinsic kinematic properties of the galaxies taking into account the thickness of the HI disc under the assumption of vertical hydrostatic equilibrium. Applying this method to WLM (Fig. 21), we derived a scale height that is about 150 pc in the centre and flares linearly with radius up to 600 pc, well above the 100 pc assumed in our analysis. We compared the results obtained taking the HI thickness into account with the results obtained in the current work as described in Sec. 4: in the thick-disc model the peak of the profile of intrinsic surface density is higher by about $2 \text{ M}_\odot \text{pc}^{-2}$, the velocity dispersion is lower by an average of 0.5 km/s and the rotation velocities have a maximum difference of about 4 km/s. All these differences are compatible within the errors and we can conclude that our results for WLM are not seriously biased by the assumption of a thin disc.

The scale height of a HI disc can be related to the velocity dispersion of the gas and the total volumetric density in the plane of the galaxy as $z_d \propto \sigma_v \rho^{-0.5}$ (Olling 1995; Ott et al. 2001), we can define an average density in terms of circular velocity so that $z_d R^{-1} \propto \sigma_v V_c^{-1}$. Therefore, WLM is expected to be representative of our sample of galaxies, in term of disc thickness (see Tab. 2): the bias due to the HI thickness is likely to be significantly larger only for DDO 210, DDO 216 and NGC 1569. However, in these galaxies the rotation curve is already very uncertain and the quoted errors are so large that should be still inclusive of the errors due to the thin-disc assumption. In conclusion, we are confident that all the rotation curves and velocity dispersions found in this work are not seriously biased by the assumption that the HI disc is thin.

7.2 Comparison to the standard 2D approach

In this Section we compare the rotation curves obtained with our method with the ones obtained in O15 for the same dataset with the classical 2D approach. As a general feature, the rotation curves of O15 reach larger radii: the median difference on the radial extension is about 700 pc and the galaxies with the largest discrepancies are: DDO 50 (3.4 kpc), DDO 87 (2.2 kpc) and NGC 2366 (1.8 kpc). DDO 216 has rotation curves of similar extensions, while our rotation curves for DDO 210 and DDO 168 are more extended by about 200 pc and 700 pc respectively. These differences are due to the different approach in the choices on the outermost radii considered in the kinematic fit. We decided to include only the part of the galaxy with clear information on the rotation motion and not dominated by the noise (see Sec. 4.1.2), as shown by the elliptical rings over plotted to the HI maps and HI velocity fields (Box C in Figs. 5-21). In contrast, O15 extended the fit as much as possible, but in some cases the outer part of the rotation curves might be extracted mainly from emission around the minor axis which have poor information on the rotational motion. The peril of this approach is that the rotation curves at large radii can be quite unreliable or show artifacts as for example in WLM (Fig. 24a) where the rotation curve

of O15 shows an abrupt drop-off after the outermost radii used in our analysis.

Concerning the shapes of rotation curves, about half of our results are compatible with O15 within the errors. When differences are present they can be due to a combination of three different reasons: (i) an intrinsic difference in the best-fit projected rotational velocities ($V_{\text{rot}} \sin i$), (ii) a difference in the best-fit i , PA or centre and (iii) a difference in the asymmetric-drift correction terms (V_A). Most of the cases are due to (ii) and (iii); only one galaxy (DDO 168) seems to have a significant discrepancy caused by the two different fitting methods. This galaxy is shown in Fig. 24c, while examples of (ii) and (iii) are shown in Figs. 24b and 24d. The galaxies showing the largest discrepancies are listed below.

- **CVnidwa.** Our final rotation curve is systematically higher than that of O15 by about 5 km/s. This difference is totally dominated by the discrepancy in the asymmetric-drift correction term. Our V_A is already significant at the inner radii, while the V_A of O15 is negligible out to 2 kpc. These differences are ascribed to the different radial trend of the intrinsic surface density profile caused by discrepancies in the best-fit PA.
- **DDO 47.** The circular velocity shows a discrepancy of about 7 km/s in the inner part of the disc ($R < 2$ kpc) due to a discrepancy on the best-fit i : O15 used $i \approx 55^\circ$ while we used $i = 37^\circ$ independent of radius.
- **DDO 50.** Discrepancies of about 10 km/s (see Fig. 24b) are due to the different assumption on the value of i (about 50° in O15 and 33° in this work). Our curve is also less extended (see Fig. 7).
- **DDO 52.** The final rotation curve of O15 is systemically higher due to the different values of i (about 37° in O15 and 55° in this work).
- **DDO 87.** Our final rotation curve shows a steeper rising due to a discrepancy in the assumed i : we used $i = 40^\circ$ independent of radius, while in O15 i varies between 60° and 40° .
- **DDO 168.** The final rotation curves are quite different (see Fig. 24c). In O15 the curve flattens at about 2 kpc and starts to decrease at 3 kpc, while our rotation curve shows a less steep rising in the inner part and a flattening at about 3.5 kpc. These discrepancies are due mainly to an intrinsic difference in the best-fit projected rotation curve found with the two methods. The differences are further increased by the different assumptions on i ($i \approx 60^\circ$ in this work, $i \approx 40^\circ$ in O15).
- **NGC 1569.** The circular rotation curve found in this work rises steeper with respect to the rotation curve reported in O15 (see Fig. 24d). The difference is caused by the asymmetric-drift correction given that with our 3D method we find a different radial trend of both the velocity dispersion and the intrinsic surface density.
- **NGC 2366.** Our final rotation curve is systematically lower in the inner disc ($R < 2$ kpc). The cause is a combination of a different position of the galactic centre and different PA. We used a constant PA of about 40° , while in O15 the PA grows from 20° to 40° in the first two kpc.

8 SUMMARY

We presented a study of the kinematics of the HI discs for 17 dwarf irregular galaxies taken from the public survey LITTLE THINGS (Hunter et al. 2012). The main goal of this work is to make available to the community a sample of high-quality rotation curves of dIrrs ready-to-use to perform dynamical studies. The tabulated quantities (V_{rot} , V_c , V_A , σ_v) and HI surface-density profile (Σ_{int})




Figure 24. Circular velocity (V_c) and asymmetric-drift term (V_A) for four representative galaxies in our sample (a: WLM, b: DDO50, c: DDO 168, d: NGC 1569). The red squares and the red curves show the results found in this work, while the grey circles and the grey curves show the results obtained in O15.

are available in the online version of the paper. The key points of this work are listed below.

1. We derived the rotation curves from the HI datacubes with a state-of-art technique: the publicly available software ^{3D}BAROLO (Di Teodoro & Fraternali 2015). It fits 3D models to the datacubes without explicitly extracting velocity fields. The fit in the 3D space of the datacubes ensures full control of the observational effects and in particular a proper account of beam smearing.
2. The maximum radii used in the fits have been chosen with great care from the HI total map. This ensures a robust estimate of the kinematics avoiding regions of the galaxy dominated by the noise and with scant information about the gas rotation.
3. We developed a method to take into account the uncertainties of the asymmetric-drift correction. As a consequence, the quoted errors on the rotation curves are representative of the real uncertainties. The inclusion of these errors are fundamental in galaxies for which the calculation of the circular velocity is highly dominated by the asymmetric-drift correction (e.g. DDO 210).
4. We estimated how our results can be biased by the assumption of a thin HI disc. We analysed in detail the galaxy WLM and we found that the HI layer flares linearly from about 150 pc to 600

pc, well above the 100 pc assumed in our analysis. The presence of such thick disc introduces systematic errors on the estimate of the velocity dispersion, intrinsic HI surface density and circular velocity. However, the differences are compatible within the quoted uncertainties. The thickness of WLM should be representative of the galaxies of our sample, so we can conclude that our results are not seriously biased by the assumption of thin disc.

5. The rotation curves obtained in this work have been used to test the baryonic Tully-Fisher relation in the low-mass regime ($10^6 \lesssim M_{\text{bar}}/M_{\odot} \lesssim 10^9$). We found that our results are compatible in slope, normalisation and scatter with the work of Lelli et al. (2016b) at higher mass ($M_{\text{bar}} > 10^8 M_{\odot}$).

Acknowledgments: This research made use of the LITTLE THINGS data sample. We would like to thank the anonymous referee and Antonino Marasco for useful comments that improved this manuscript and Se-Heon Oh and Federico Lelli for kindly making their data available. G. Battaglia acknowledges financial support by the Spanish Ministry of Economy and Competitiveness (MINECO) under the Ramon y Cajal Programme (RYC-2012-11537). J.I. Read

would like to acknowledge support from STFC consolidated grant ST/M000990/1 and the MERAC foundation. The research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

REFERENCES

- Baillard A., et al., 2011, *A&A*, **532**, A74
- Battaglia G., Fraternali F., Oosterloo T., Sancisi R., 2006, *A&A*, **447**, 49
- Begeman K. G., 1987, PhD thesis, , Kapteyn Institute, (1987)
- Begum A., Chengalur J. N., Karachentsev I. D., Sharina M. E., 2008, *MNRAS*, **386**, 138
- Bosma A., 1978a, PhD thesis, PhD Thesis, Groningen Univ., (1978)
- Bosma A., 1978b, PhD thesis, PhD Thesis, Groningen Univ., (1978)
- Bosma A., 1981, *AJ*, **86**, 1791
- Bosma A., Goss W. M., Allen R. J., 1981, *A&A*, **93**, 106
- Boylan-Kolchin M., Bullock J. S., Kaplinghat M., 2011, *MNRAS*, **415**, L40
- Brinks E., Burton W. B., 1984, *A&A*, **141**, 195
- Brook C. B., Santos-Santos I., Stinson G., 2016a, *MNRAS*, **459**, 638
- Brook C. B., Santos-Santos I., Stinson G., 2016b, *MNRAS*, **459**, 638
- Bureau M., Carignan C., 2002, *AJ*, **123**, 1316
- Carignan C., Purton C., 1998, *ApJ*, **506**, 125
- Casertano S., van Gorkom J. H., 1991, *AJ*, **101**, 1231
- Cole A. A., et al., 1999, *AJ*, **118**, 1657
- Cook D. O., et al., 2014, *MNRAS*, **445**, 881
- Côté S., Carignan C., Freeman K. C., 2000, *AJ*, **120**, 3027
- Croxall K. V., van Zee L., Lee H., Skillman E. D., Lee J. C., Côté S., Kennicutt Jr. R. C., Miller B. W., 2009, *ApJ*, **705**, 723
- Di Cintio A., Lelli F., 2016, *MNRAS*, **456**, L127
- Di Teodoro E. M., Fraternali F., 2015, *MNRAS*, **451**, 3021
- Fraternali F., Oosterloo T., Sancisi R., van Moorsel G., 2001, *ApJ*, **562**, L47
- Fraternali F., van Moorsel G., Sancisi R., Oosterloo T., 2002, *AJ*, **123**, 3124
- García-Ruiz I., Sancisi R., Kuijken K., 2002, *A&A*, **394**, 769
- Gentile G., Salucci P., Klein U., Vergani D., Kalberla P., 2004, *MNRAS*, **351**, 903
- Gentile G., Burkert A., Salucci P., Klein U., Walter F., 2005, *ApJ*, **634**, L145
- Georgiev T. B., Karachentsev I. D., Tikhonov N. A., 1997, *Astronomy Letters*, **23**, 514
- Hunter D. A., Elmegreen B. G., 2006, *ApJS*, **162**, 49
- Hunter D. A., Elmegreen B. G., van Woerden H., 2001, *ApJ*, **556**, 773
- Hunter D. A., Elmegreen B. G., Oh S.-H., Anderson E., Nordgren T. E., Massey P., Wilsey N., Riabokin M., 2011, *AJ*, **142**, 121
- Hunter D. A., et al., 2012, *AJ*, **144**, 134
- Jarrett T. H., Chester T., Cutri R., Schneider S. E., Huchra J. P., 2003, *AJ*, **125**, 525
- Johnson M., Hunter D. A., Oh S.-H., Zhang H.-X., Elmegreen B., Brinks E., Tollerud E., Herrmann K., 2012, *AJ*, **144**, 152
- Józsa G. I. G., Kenn F., Klein U., Oosterloo T. A., 2007, *A&A*, **468**, 731
- Kamphuis P., Józsa G. I. G., Oh S.-H., Spekkens K., Urbancic N., Serra P., Koribalski B. S., Dettmar R.-J., 2015, *MNRAS*, **452**, 3139
- Karachentsev I. D., Makarov D. I., Kaisina E. I., 2013, *AJ*, **145**, 101
- Kirby E. N., Bullock J. S., Boylan-Kolchin M., Kaplinghat M., Cohen J. G., 2014, *MNRAS*, **439**, 1015
- Klypin A., Kravtsov A. V., Valenzuela O., Prada F., 1999, *ApJ*, **522**, 82
- Knapen J. H., Erroz-Ferrer S., Roa J., Bakos J., Cisternas M., Leaman R., Szymanek N., 2014, *A&A*, **569**, A91
- Krajnović D., Cappellari M., de Zeeuw P. T., Copin Y., 2006, *MNRAS*, **366**, 787
- Leaman R., et al., 2012, *ApJ*, **750**, 33
- Lelli F., Fraternali F., Sancisi R., 2010, *A&A*, **516**, A11
- Lelli F., Verheijen M., Fraternali F., Sancisi R., 2012a, *A&A*, **537**, A72
- Lelli F., Verheijen M., Fraternali F., Sancisi R., 2012b, *A&A*, **544**, A145
- Lelli F., Verheijen M., Fraternali F., 2014a, *MNRAS*, **445**, 1694
- Lelli F., Verheijen M., Fraternali F., 2014b, *A&A*, **566**, A71
- Lelli F., McGaugh S. S., Schombert J. M., 2016a, *ApJ*, **816**, L14
- Lelli F., McGaugh S. S., Schombert J. M., 2016b, *ApJ*, **816**, L14
- Makarova L., 1999, *A&AS*, **139**, 491
- McConnachie A. W., 2012, *AJ*, **144**, 4
- McGaugh S. S., 2012, *AJ*, **143**, 40
- McGaugh S. S., Schombert J. M., Bothun G. D., de Blok W. J. G., 2000, *ApJ*, **533**, L99
- O’Brien J. C., Freeman K. C., van der Kruit P. C., 2010, *A&A*, **515**, A62
- Oh S.-H., de Blok W. J. G., Walter F., Brinks E., Kennicutt Jr. R. C., 2008, *AJ*, **136**, 2761
- Oh S.-H., Brook C., Governato F., Brinks E., Mayer L., de Blok W. J. G., Brooks A., Walter F., 2011, *AJ*, **142**, 24
- Oh S.-H., et al., 2015, *AJ*, **149**, 180
- Olling R. P., 1995, *AJ*, **110**, 591
- Olling R. P., 1996, *AJ*, **112**, 481
- Oman K. A., Navarro J. F., Sales L. V., Fattahi A., Frenk C. S., Sawala T., Schaller M., White S. D. M., 2016, preprint, ([arXiv:1601.01026](https://arxiv.org/abs/1601.01026))
- Ott J., Walter F., Brinks E., Van Dyk S. D., Dirsch B., Klein U., 2001, *AJ*, **122**, 3070
- Puche D., Westpfahl D., Brinks E., Roy J.-R., 1992, *AJ*, **103**, 1841
- Read J. I., Wilkinson M. I., Evans N. W., Gilmore G., Kleyna J. T., 2006, *MNRAS*, **367**, 387
- Read J. I., Iorio G., Agertz O., Fraternali F., 2016a, preprint, ([arXiv:1607.03127](https://arxiv.org/abs/1607.03127))
- Read J. I., Agertz O., Collins M. L. M., 2016b, *MNRAS*, **459**, 2573
- Read J. I., Iorio G., Agertz O., Fraternali F., 2016c, *MNRAS*, **462**, 3628
- Roberts M. S., 1975, *Radio Observations of Neutral Hydrogen in Galaxies*. the University of Chicago Press, p. 309
- Rogstad D. H., Shostak G. S., 1971, *A&A*, **13**, 99
- Rogstad D. H., Lockhart I. A., Wright M. C. H., 1974, *ApJ*, **193**, 309
- Roychowdhury S., Chengalur J. N., Begum A., Karachentsev I. D., 2010, *MNRAS*, **404**, L60
- Schoenmakers R. H. M., Franx M., de Zeeuw P. T., 1997, *MNRAS*, **292**, 349
- Sicking F. J., 1997, PhD thesis, PhD Thesis, University of Groningen, (1997)
- Spekkens K., Sellwood J. A., 2007, *ApJ*, **664**, 204
- Stil J. M., Israel F. P., 2002a, *A&A*, **389**, 42
- Stil J. M., Israel F. P., 2002b, *A&A*, **392**, 473
- Swaters R. A., 1999, PhD thesis, , Rijksuniversiteit Groningen, (1999)
- Swaters R. A., Schoenmakers R. H. M., Sancisi R., van Albada T. S., 1999, *MNRAS*, **304**, 330
- Tamburro D., Rix H.-W., Leroy A. K., Mac Low M.-M., Walter F., Kennicutt R. C., Brinks E., de Blok W. J. G., 2009, *AJ*, **137**, 4424
- Taylor C. L., Kobulnicky H. A., Skillman E. D., 1998, *AJ*, **116**, 2746
- Verheijen M. A. W., 2001, *ApJ*, **563**, 694
- Verheijen M. A. W., Sancisi R., 2001, *A&A*, **370**, 765
- Walter F., Brinks E., 2001, *AJ*, **121**, 3026
- Whiting M. T., 2012, *MNRAS*, **421**, 3242
- Zhang H.-X., Hunter D. A., Elmegreen B. G., Gao Y., Schruba A., 2012, *AJ*, **143**, 47
- de Blok W. J. G., Bosma A., 2002a, *A&A*, **385**, 816
- de Blok W. J. G., Bosma A., 2002b, *A&A*, **385**, 816
- de Blok W. J. G., Walter F., Brinks E., Trachternach C., Oh S.-H., Kennicutt Jr. R. C., 2008, *AJ*, **136**, 2648
- van Albada T. S., Bahcall J. N., Begeman K., Sancisi R., 1985, *ApJ*, **295**, 305
- van Eymeren J., Marcelin M., Koribalski B., Dettmar R.-J., Bomans D. J., Gach J.-L., Balard P., 2009a, *A&A*, **493**, 511
- van Eymeren J., Marcelin M., Koribalski B. S., Dettmar R.-J., Bomans D. J., Gach J.-L., Balard P., 2009b, *A&A*, **505**, 105
- van der Hulst J. M., Terlouw J. P., Begeman K. G., Zwitter W., Roelfsema P. R., 1992, in Worrall D. M., Biemesderfer C., Barnes J., eds, *Astronomical Society of the Pacific Conference Series Vol. 25, Astronomical Data Analysis Software and Systems I*. p. 131




Figure A1. Initial estimate of the inclination angle for DDO 52 with 3DB (left-hand panel) and with the fit of the iso-density contours of the HI total map (right-hand panel). The red lines show the inclination of the optical disc. See the text for details.




Figure A2. Same as Fig. A1 but for DDO 210.

APPENDIX A: ESTIMATE OF THE GEOMETRICAL PARAMETERS

It is important to initialise the fit of the HI datacubes with good estimates of the geometrical parameters: centre, i and PA of the HI disc (see Sec. 4.2). We used three different methods:

- **3DB:** 3DB can automatically estimate the geometrical parameters analysing the 0th moment (HI total map) and the 1th moment (velocity field) of the datacube. The i is estimated by fitting HI model maps to the observed HI total map, while the PA is calculated as the orientation of the line that maximises the velocity gradient in the velocity field. Finally, the centre is estimated as the flux-weighted average position of the source. 3DB builds the total map and the velocity field using a source-finding algorithm (DUCHAMP, Whiting 2012), separating the source pixels from the noise pixels on the basis of a S/N threshold. As a consequence, the final maps and the estimate of the disc parameters depend on the choice of this threshold. For this reason we repeat this analysis using S/N threshold ranging from 2.5 to 5. Finally, we average the values on the range of S/N where the estimated parameters are almost constant.
- **RingFit:** We developed a simple script, called RingFit, to fit ellipses to the iso-density contours at different radii of our total HI maps (Sec. 4.1.1). From the properties of the fitted ellipses we can estimate the geometrical parameters of the HI disc at different radii.
- **Isophotal fitting of the optical maps:** The geometrical parameters are derived from the V-Band photometry by Hunter & Elmegreen (2006).

Figs. A1 and A2 show two examples of the estimate of i with the three different methods. In the first case (DDO 52) the methods are in good agreement, while in the second case (DDO 210) the results are not compatible (notice that in this case the HI contours deviate strongly from elliptical shape, see Box C in Fig. 16). Usually, the optical disc has a more regular morphology with respect to the HI disc, therefore we give more weight to the geometrical parameters estimated from Hunter & Elmegreen (2006). When the geometry of the stellar and of the HI disc do not show a clear agreement (Fig. A2) or the stellar disc presents peculiar features such as strong bars, we use the other two methods. More details can be found in the description of the individual galaxies in Sec. 5.

APPENDIX B: PLOT LAYOUT

In this Appendix we report the caption that describes Figs. 5-21.

- **Box A:** Rotation velocity estimated with 3DB (V_{rot} , blue line), asymmetric-drift correction (V_A , grey circles) and final corrected circular velocity (V_c , black circles).

- **Box B:**

Top: Inclination (left-hand panel) and position angle (right-hand panel) found with 3DB as functions of radius R . The red lines show the fit used to obtain the rotation velocity (see Sec. 4.2), while the black dashed line indicates the median of the data.

Bottom: HI velocity dispersion estimated with 3DB (left-hand panel) and intrinsic surface density of the HI disc not corrected for Helium (right-hand panel) as function of radius R . In the left-hand panel the black dashed line indicates the median of the data. The empty circles (if present) represent the points excluded both from the calculation of the asymmetric-drift correction (see Sec. 4.3.1) and from the calculation of the median. The excluded points belong to rings where the velocity dispersion found with 3DB is discrepant or peculiar with respect to the global trend (see Sec. 4.3.4).

- **Box C:**

Left-hand panel: Stellar emission overlaid with the iso-density contours of the total HI emission. The thick contour indicates σ_{3T} , the $3\text{-}\sigma$ pseudo noise of the total map (see Sec. 4.1.2 and Tab. 1).

Right-hand panel: Velocity field obtained as the 1th moment of the data cube. The thick contour highlights the systemic velocity (see Tab. 2). A physical scale is plotted on the bottom right corner of the panel, while the beam of the HI observation is shown in the bottom left corner.

The crosses and the ellipses show the assumed galactic centre and the last used ring respectively (see Tab. 2).

- **Box D:** PV diagram along the major (left-handed panel) and the minor (right-handed panel) axis of the HI disc. The black and the red contours show respectively the iso-density contours of the galaxy and the best-fit model found with 3DB. The horizontal black dashed lines indicate the systemic velocity.

APPENDIX C: AN ALTERNATIVE SCENARIO FOR DDO 216

DDO 216 shows a peculiar kinematics: a velocity gradient of about 10 km/s appears clearly in the velocity field (right-hand panel in Box C in Fig. 17) and in the channel maps, but it could be entirely due to a single ‘cloud’ at a discrepant velocity in the approaching side of the galaxy (Stil & Israel 2002a). The analysis of the channel maps partially supports this hypothesis: the receding and the approaching sides of the galaxy seem to have a slightly distinct kinematics overlapping between -186 km/s and -190 km/s, while the V_{sys} found with 3DB is around -180 km/s. At the same time, the total HI map does not show the clear presence of a separate component at the supposed position of the cloud (left-hand panel in Box C in Fig. 17). In conclusion we did not have sufficient elements to confirm or exclude the existence of a HI component other than the disc of DDO 216. We can summarise the two hypothesis as:

- *scenario1*: The main velocity gradient is genuine and totally due to the rotation of the HI disc.
- *scenario2*: The main velocity gradient is due to an HI substructure with a systemic velocity that differs of about 20 km/s with respect to the HI disc.

The scenario1 is analysed in Sec. 5 while in this Appendix we describe the analysis of the scenario2. First of all we masked the region of the datacube containing the emission of the supposed HI cloud. The mask has been build by inspecting the channel maps using the task PYBLOT of GIPSY. Then we performed the kinematic fit of the ‘clean’ datacube with 3DB. We set i_{ini} , PA_{ini} and the galactic centre to the values obtained from the stellar disc (Hunter & Elmegreen 2006). Contrarily to the scenario1 (Sec. 5) we used the V_{sys} estimated with 3DB, the two estimates differ by about 8 km/s, while the final best-fit PA (120°) is tilted of about 10° - 15° with respect to the values found in the scenario1 (Box B in Fig. 17).

Without the ‘cloud’ the velocity gradient almost disappears and the kinematics is dominated by the chaotic motion of the gas as it is clearly visible in the PVs in Fig. C1. As a consequence, the final rotation curve is ‘shaped’ by the asymmetric-drift correction. Fig. C2 compares the rotation curve found for the scenario1 and for the scenario2: the velocities are compatible within the errors, but in scenario1 the V_c is systematically higher by about 4 km/s and it extends to larger radii.

Kirby et al. (2014) found that the stellar component of DDO 216 is rotating in the same direction as the gas with a V_{rot} of about 10 km/s, perfectly compatible with scenario1 (see Fig. C2). This velocity gradient is observed also in red giant stars, so the rotation is genuine and it is not caused by short-term hydrodynamical events (e.g. bubbles), which influence the gas kinematics but not the kinematics of the old stellar populations. However the centre of the rotation and V_{sys} are quite different from what we found for scenario1 and they are instead compatible with scenario2. Further studies are needed to understand if the gas in this peculiar galaxy is rotating, which is crucial to accurately estimate its dynamical mass.

APPENDIX D: TYPICAL SETUP FOR 3^{D} BAROLO

In this Appendix we report the keywords and their values for our typical parameter file passed to 3DB. For a detailed description of




Figure C1. PV along the major axis (upper panel) and the minor axis (bottom panel) for the scenario2 of the galaxy DDO 216 (see text). The black contours indicate the datacube while the red contours show the best-fit model found with 3DB. Contours at $(2 + 6n)\sigma_{\text{ch}}$, where $\sigma_{\text{ch}} = 0.91$ mJy bm^{-1} , the grey contours are at $-2\sigma_{\text{ch}}$. The yellow points indicate the rotational velocity found with 3DB.




Figure C2. Rotational velocities (red dashed line) and circular velocities (red points) found for the scenario2 of DDO 216 (see text). The rotation curves of the scenario1 are also shown for comparison.

the keywords see the documentation of 3DB⁶ ('...' means that the values depends on the analysed galaxy).

- FITSFILE ...; name of the datacube to fit.
- GALFIT TRUE; enable the 3D fitting of the datacube.
- NRADII ...; number of rings to be used in the fit.
- RADSEP ...; separation between rings in arcsec.
- VROT 30; initial guess for the rotation velocity in km/s.
- VDISP 8; initial guess for the velocity dispersion in km/s.
- INC ...; initial guess for i in degree (i_{ini} , see Tab. 2).
- PA ...; initial guess for the PA in degree (PA_{ini} , see Tab. 2).
- VSYS ...; initial guess for the systemic velocity in km/s.
- XPOS ...; initial guess for the X coordinate of the centre.
- YPOS ...; initial guess for the Y coordinate of the centre.
- Z0 $100/f_c$; initial guess for the scale height in arcsec, f_c is the conversion factor from arcsec to pc and it is listed in Tab.1 for each galaxy.
- LTYPE 1; use a Gaussian profile for the HI vertical distribution.
- FREE VROT VDISP INC PA; parameters to fit with 3DB.
- FTYPE 2; use the absolute differences between models and datacube as residuals.
- WFUNC 1; use the cosine of the azimuthal angle as weighing function for the fit (see Sec. 4.2).
- CDENS 10; this parameter sets (in a non-trivial way) the number of 'clouds' to be used in the building of the ring models.
- NV 200; number of sub-clouds in which each cloud (see above) is divided to populate the spectral axis of the ring models.
- TOL 1E-3; tolerance condition to stop the fit.
- MASK Smooth; create a mask for the fit smoothing the datacube.
- BLANKCUT 2.5; include in the mask (see above) only the pixels with $S/N > 2.5$.
- SIDE B; fit the whole galaxy.
- TWOSTAGE TRUE; enable the two-stage method described in Sec. 4.2.
- BWEIGHT 1; set the weight of the blank pixels (see Di Teodoro & Fraternali 2015 for details).
- DELTAINC 40; i can assume values only between INC-DELTAINC and INC+DELTAINC.
- DELTAPA 80; same as above, but for the PA.
- FLAGERRORS true; enable the estimate of the errors on the fitted parameters.
- STARTRAD 1; avoid the fit of the first ring placed by default at $\text{RADSEP}/4$.
- LINEAR 0.424; instrumental resolution (σ_{inst}) in unity of channel width. In all our galaxies the instrumental FWHM is equal to the channel separation, so $\sigma_{\text{inst}} = 1/2.355 = 0.424$.

If the parameters XPOS,YPOS,VSYS,PA,INC are omitted, 3DB estimates them as described in Appendix A.

This paper has been typeset from a \LaTeX file prepared by the author.

⁶ <http://editeodoro.github.io/Bbarolo/documentation/>